

Operator mixing in $\mathcal{N} = 4$ SYM: The Konishi anomaly re-re-visited

B. Eden

*Institut für theoretische Physik, Universität Leipzig
Postfach 100920, D-04009 Leipzig, Germany
burkhard.eden@itp.uni-leipzig.de*

Abstract:

The supersymmetry transformation relating the Konishi operator to its lowest descendant in the 10 of $SU(4)$ is not manifest in the $\mathcal{N} = 1$ formulation of the theory but rather uses an equation of motion. On the classical level one finds one operator, the unintegrated chiral superpotential. In the quantum theory this term receives an admixture by a second operator, the Yang-Mills part of the Lagrangian. It has long been debated whether this “anomalous” contribution is affected by higher loop corrections. We present a first principles calculation at the second non-trivial order in perturbation theory using supersymmetric dimensional reduction as a regulator and renormalisation by Z-factors. Singular higher loop corrections to the renormalisation factor of the Yang-Mills term are required if the conformal properties of two-point functions are to be met. These singularities take the form determined in preceding work on rather general grounds. Moreover, we also find non-vanishing finite terms.

The core part of the problem is the evaluation of a four-loop two-point correlator which is accomplished by the Laporta algorithm. Apart from several examples of the T_1 topology with two lines of non-integer dimension we need the first few orders in the ϵ expansion of three master integrals. The approach is self-contained in that all the necessary information can be derived from the power counting finiteness of some integrals.

1 Introduction

In the $\mathcal{N} = 4$ super Yang-Mills theory ($\mathcal{N} = 4$ SYM), half BPS operators are known to be protected. Their two- and three-point functions receive no quantum corrections other than contact terms [1, 2]. The simplest such operator is

$$\mathcal{O} = \text{Tr}(\phi^1 \phi^1) \quad (1)$$

where the notation refers to the action of the theory (91) in terms of $\mathcal{N} = 1$ superfields [3]. The physical fields are the $\theta = \bar{\theta} = 0$ components of the three complex chiral fields and of

$$\begin{aligned} \lambda_\alpha^1 &= \lambda_\alpha &= \frac{1}{4g} \bar{D}_{\dot{\alpha}} \bar{D}^{\dot{\alpha}} (e^{gV} D_\alpha e^{-gV}) \\ \lambda_\alpha^i &= \nabla_\alpha \phi^{i-1} &= e^{gV} (D_\alpha (e^{-gV} \phi^{i-1} e^{gV})) e^{-gV}, \quad i \in \{2, 3, 4\} \end{aligned} \quad (2)$$

and their complex conjugates, while the gauge field is in the $[D_\alpha, \bar{D}_\alpha]$ component of V . The little $i \in \{1 \dots 4\}$ is an index in the fundamental representation of the non-manifest but unbroken $SU(4)$ R-symmetry of the theory. Using the equation of motion

$$-\frac{1}{4} D^\alpha D_\alpha \phi^I = \frac{g}{2} \varepsilon^{IJK} [\bar{\phi}_J, \bar{\phi}_K], \quad I \in \{1, 2, 3\} \quad (3)$$

of the chiral field we find

$$\begin{aligned} \frac{1}{2} D^\alpha D_\alpha \mathcal{O} &= \check{F} - 4g \check{B}, \\ \check{F} &= \text{Tr}(\lambda^{2\alpha} \lambda_\alpha^2), \quad \check{B} = \text{Tr}((e^{-gV} \phi^1 e^{gV}) [\bar{\phi}_2, \bar{\phi}_3]). \end{aligned} \quad (4)$$

The operator $\check{F}$ is an $SU(4)$ component of

$$F^{10,ij} = \text{Tr}(\lambda^{i\alpha} \lambda_\alpha^j), \quad (5)$$

and similarly for $\check{B}$. The associated highest weight components are

$$F = \text{Tr}(\lambda^\alpha \lambda_\alpha), \quad B = \text{Tr}(\phi^1 [\phi^2, \phi^3]). \quad (6)$$

We verify by explicit graph calculations at $O(g^2)$ in Section 3 for the highest weight components, and at $O(g^4)$ in Section 5 for the 22 components $\check{F}$, $\check{B}$, that the combination

$$O^{10} = F^{10} - 4g B^{10} \quad (7)$$

is a protected operator. Further, the contact contributions in its two-point function are related to those of the $\mathcal{O}$ case by the classical e.o.m. (3), see Section 5.

Second, from the complex conjugate of the classical relation (3) we find

$$-\frac{1}{4} \bar{D}_{\dot{\alpha}} \bar{D}^{\dot{\alpha}} \mathcal{K} = -3g B, \quad \mathcal{K} = \text{Tr}(e^{gV} \bar{\phi}_I e^{-gV} \phi^I). \quad (8)$$

The operator $\mathcal{K}$ (the kinetic term for matter) is usually called the “Konishi operator”.

The conformal symmetry of $\mathcal{N} = 4$ SYM implies that two-point functions of properly renormalised operators must take the form of a power law (N is the rank of the gauge group $SU(N)$)

$$\langle \bar{O}_r(x_2) O_r(x_1) \rangle = \frac{c_r(g^2, N)}{(x_{12}^2)^{\Delta_r(g^2, N)}}, \quad x_{12} = x_1 - x_2; \quad \langle \bar{O}_r(x_2) O_s(x_1) \rangle = 0 : \quad \Delta_r \neq \Delta_s \quad (9)$$

where r, s labels various operators. Quantum corrections can thus affect the normalisation c or lead to logarithms that sum into the “anomalous dimension”, i.e. the g^2 dependent part of Δ . Both equations only hold up to contact terms; in dimensional regularisation or related schemes this means terms of order ϵ before the regulator is sent to zero. Both $\mathcal{O}$ and O^{10} trivially satisfy the first equation in (9) because quantum corrections are absent, in particular $\Delta = 2, 3$, respectively. On the other hand, $\mathcal{K}$ must be renormalised. By various arguments (first from the OPE [4]) it is known that its dimension behaves like

$$\Delta_{\mathcal{K}} = 2 + 3\alpha - 3\alpha^2 + \dots, \quad \alpha = \frac{g^2 N}{4\pi^2}. \quad (10)$$

Curiously, B as the “descendant” of $\mathcal{K}$ under the classical equation of motion is not orthogonal to $O = F - 4gB$. Moreover, from the explicit results below we would calculate an anomalous dimension $\Delta_B = 3 + 3\alpha - 9/2\alpha^2 + \dots$ and hence not the same as for $\mathcal{K}$. This is in contradiction to supersymmetry: The $\mathcal{N} = 1$ supersymmetric version of the power law in (9) is obtained by replacing x_{12} by a supersymmetric line element. Applying the differential operator from the equation of motion on either end (thus $\bar{D}^2|_1 D^2|_2$) simply produces a box operator (we illustrate this in Section 5 on the simpler case $\mathcal{O}$). The dimension of the correct descendant is thus higher by one unit whereas its anomalous part must agree.

According to [5, 6] in the case of $\mathcal{K}$ the equation of motion is modified in the quantum theory: The correct descendant is

$$K := \frac{1}{12g} \bar{D}_{\dot{\alpha}} \bar{D}^{\dot{\alpha}} \mathcal{K} = B + \frac{gN}{32\pi^2} F + \dots \quad (11)$$

This is an effect of the renormalisation of composite operators and not a supersymmetry anomaly. When $\mathcal{K}$ is regularised by point splitting it is in fact possible to derive the lowest order F admixture from the supersymmetry variation of a Wilson line between the separated chiral fields in $\mathcal{K}$ [6]. The mixture K is orthogonal to O at order g^1 , and we show in the present article that it has the same anomalous dimension as $\mathcal{K}$ up to $O(g^4)$.

In theories without matter self-interaction the classical equation of motion would send $\mathcal{K}$ to zero, which implies the conservation of an axial current in the $\mathcal{N} = 1$ Grassmann expansion of the operator. In the literature, the F admixture therefore has been coined the “Konishi anomaly” in analogy to the standard axial anomaly. In several models without superpotential the F term does in fact not receive higher loop corrections [7]. The more general case with matter self-interaction is included in a similar statement in [8].

In [9] the question was taken up again in $\mathcal{N} = 4$ SYM by an analysis of the conformal properties of two-point functions as stated in (9). Unfortunately, point splitting is awkward to use at higher orders in perturbation theory. Therefore the discussion was build around supersymmetric dimensional reduction (SSDR) as a regulator [10], on the expense of losing the direct derivation of the “anomaly”. On rather general grounds (the singularity structure and fractional dimension of the correlators, supplemented by parametric differentiation as in renormalisation group reasoning) it was shown that the “anomaly” *must* be affected by renormalisation at least in the given scheme, in apparent disagreement with [8].

To lend further support to the claim, we present a perturbative treatment in SSDR at the second non-trivial order.¹ As in [9] we resort to the conformal properties (9) to resolve the mixing. The programme is carried out from first principles: It is shown that the two-point function of $O = F - 4gB$ is protected at $O(g^2)$ and $O(g^4)$. We then impose orthogonality of

$$K = Z_B B + \frac{gN}{32\pi^2} Z_F F \quad (12)$$

to O and the conformal form of the two-point function of K at the first two non-trivial orders in g^2 . Of necessity, Z_F contains singular higher loop corrections.

Upon rewriting $F = O + 4gB$ and rescaling, the “Konishi descendant” takes the form

$$\tilde{K} = \tilde{Z}_B B + \frac{gN}{32\pi^2} \tilde{Z}_O O \quad (13)$$

for which the Z-factors have been worked out in [9] in closed form in terms of the anomalous part of the dimension:

$$\begin{aligned} \tilde{Z}_B &= \exp\left(\frac{1}{2\epsilon} \int_0^\alpha \frac{\gamma(\tau)}{\tau} d\tau\right), \quad \gamma(\tau) = \gamma_1 \tau + \gamma_2 \tau^2 + \dots \\ \tilde{Z}_O &= \frac{1}{\alpha} \left(\omega(\alpha) - \tilde{Z}_B(\alpha, \epsilon) \int_0^\alpha \omega(\tau) \partial_\tau \left(\frac{1}{\tilde{Z}_B(\tau, \epsilon)} \right) d\tau \right), \quad \omega(\tau) = \tau + \omega_1 \tau + \omega_2 \tau^2 + \dots \end{aligned} \quad (14)$$

Note that the two Z-factors are not proportional. Apart from the anomalous dimension $\gamma(\alpha)$, graph calculations can only determine the finite part $\omega(\alpha)$ of $\tilde{Z}_O$. In terms of this data the poles in both Z-factors are given by the formulae above. In particular, $\tilde{Z}_O$ is singular even if $\omega_i = 0$.

The value $\omega_1 = -3/4$ was given in [9], below we establish $\omega_2 = 7/8$. Moreover, up to $O(g^4)$ the singular terms in the renormalisation factors obtained by direct calculation coincide with the values expected from (14).

¹Potential ambiguities in SSDR at very high orders [11] do not play a role in this work.

2 Loop corrections to the Konishi anomaly in SSDR

In the next three sections of the article we work out the correlators $\langle \bar{B}B \rangle$, $\langle \bar{F}F \rangle$ and $\langle \bar{B}F \rangle$ at the first three orders in perturbation theory. We use $\mathcal{N} = 1$ superfields in supersymmetric Fermi-Feynman gauge. The regulator is supersymmetric dimensional reduction. The hard part of the problem is $\langle \bar{B}F \rangle_{g^5}$ which contains 58 four-loop two-point superdiagrams. Breaking down the Grassmann algebra does not lead to simple numerators. We have evaluated the integrals on the computer using the Laporta algorithm, i.e. integration by parts. A sketch of the approach is given in Section 6. The numerator reduction has been independently verified by A. Pak.

In this section we discuss the operator mixing problem between B and F drawing upon the results, without touching upon the details of the graph calculation. The renormalisation procedure laid out below is insensitive to division by regular functions of ϵ , the decrement of the space time dimension $D = 4 - 2\epsilon$. In equation (17) we have scaled down the correlators by $\langle \bar{B}B \rangle_{g^0}$, so

$$BB_2 = \frac{\langle \bar{B}(x_2)B(x_1) \rangle_{g^2}}{\langle \bar{B}(x_2)B(x_1) \rangle_{g^0}} \quad (15)$$

etc. and we use the abbreviations (at the given orders there are no corrections subleading in N)

$$X = x_{12}^2 \tilde{\mu}_x^2, \quad \alpha = \frac{g^2 N}{4\pi^2}. \quad (16)$$

The vanishing of $\langle \bar{B}F \rangle_{g^1}$ is a trivial consequence of the Feynman rules; there is no diagram.

$$\begin{aligned} BB_0 &= 1 \\ BB_2 &= \frac{\alpha X^\epsilon}{\epsilon} \left[-3 - 3\epsilon - 9\zeta(3)\epsilon^2 + \dots \right] \\ BB_4 &= \left(\frac{\alpha X^\epsilon}{\epsilon} \right)^2 \left[\frac{9}{2} + \frac{45}{4}\epsilon + \left(\frac{45}{4} + \frac{63}{2}\zeta(3) \right) \epsilon^2 + \dots \right] \\ FF_0 &= 16g^2 \frac{1}{\alpha X^\epsilon} \left[2 - 4\epsilon + 2\epsilon^2 + \dots \right] \\ FF_2 &= 16g^2 \left[-1 - 12\zeta(3)\epsilon - \left(\frac{\pi^4}{5} - 54\zeta(3) \right) \epsilon^2 + \dots \right] \\ FF_4 &= 16g^2 \frac{\alpha X^\epsilon}{\epsilon} \left[-\frac{3}{2} + \frac{3}{2}\epsilon + \left(12 + 9\zeta(3) + \frac{75}{2}\zeta(5) \right) \epsilon^2 + \dots \right] \\ BF_1 &= 0 \\ BF_3 &= g \frac{\alpha X^\epsilon}{\epsilon} \left[-9 - 3\epsilon + 24\epsilon^2 + \dots \right] \\ BF_5 &= g \left(\frac{\alpha X^\epsilon}{\epsilon} \right)^2 \left[\frac{33}{2} + 35\epsilon + (-3 + 90\zeta(3)) \epsilon^2 + \dots \right] \end{aligned} \quad (17)$$

It follows immediately that the two-point function of $O = F - 4gB$ is of order ϵ at g^2 and at g^4 . In the limit $\epsilon \rightarrow 0$ these are contact terms [2]. When $x_{12} \neq 0$ the one- and two-loop corrections tend to zero — in other words, this linear combination is protected. On the other hand, protectedness at the first two orders strongly point towards all-loops protectedness. Turning the argument around we might impose protectedness and view the absence of the pole- and finite parts of the two-point function of $F - 4gB$ as a constraint relating the ϵ expansion of, say, $\langle \bar{F}F \rangle$ to that of the other correlators. In this way the leading term in FF_2 and the two leading orders in FF_4 must take the values given in the table. Moreover, FF_2 must not have a simple pole and FF_4 must not have a double pole. A graph calculation merely yields a consistency check.

The mixture O has vanishing and hence well-defined anomalous dimension. Clearly we can construct a second operator K starting on B that will have to be orthogonal to O if it has non-vanishing anomalous dimension. To lowest order in g

$$K = B + \frac{gN}{32\pi^2} F \quad (18)$$

satisfies this constraint, but we also see that

$$\langle \bar{K}(2) O(1) \rangle_{g^1} = \langle \bar{B}(2) B(1) \rangle_{g^0} 4g\epsilon \log(x_{12}^2 \tilde{\mu}_X^2) + \dots \quad (19)$$

from which it is clear that we cannot expect the orthogonality constraint to hold beyond $O(\epsilon^0)$. The bare two-point function of K is divergent at $O(g^2)$ because the pole in BB_2 cannot be compensated by the finite contribution FF_0 . We therefore renormalise as

$$\begin{aligned} K &= Z_B B + \frac{gN}{32\pi^2} Z_F F, \\ Z_B &= 1 + \alpha \frac{b_{11}}{\epsilon} + \alpha^2 \left(\frac{b_{22}}{\epsilon^2} + \frac{b_{21}}{\epsilon} \right) + \dots, \\ Z_F &= 1 + \alpha \left(\frac{f_{11}}{\epsilon} + f_{10} \right) + \alpha^2 \left(\frac{f_{22}}{\epsilon^2} + \frac{f_{21}}{\epsilon} + f_{20} \right) + \dots \end{aligned} \quad (20)$$

Finite terms in Z_B could be absorbed into an overall rescaling and a corresponding change in Z_F ; omitting these amounts to fixing the normalisation.

$$\begin{aligned} \frac{\langle \bar{K}(2) K(1) \rangle_{g^2}}{\langle \bar{B}(2) B(1) \rangle_{g^0}} &= 2\alpha \frac{b_{11}}{\epsilon} + BB_2 + \left(\frac{gN}{32\pi^2} \right)^2 FF_0 \\ &= \alpha \left[\left(2 \frac{b_{11}}{\epsilon} \right) + \left(-\frac{3}{\epsilon} - 3 - 3\log(X) \right) + \left(\frac{1}{2} \right) + O(\epsilon) \right] \end{aligned} \quad (21)$$

On the other hand, in the renormalised QFT conformal invariance implies that the two-point function has the functional form

$$\frac{\langle \bar{K}(2) K(1) \rangle_{g^2}}{\langle \bar{K}(2) K(1) \rangle_{g^0}} = \frac{1 + \alpha a_1 + \alpha^2 a_2 + \dots}{X^{\alpha\gamma_1 + \alpha^2\gamma_2 + \dots}} = 1 + \alpha(a_1 - \gamma_1 \log(X)) + O(\alpha^2). \quad (22)$$

In the first equation we can put ϵ to zero after adjusting b_{11} . It is then possible to equate the last two lines. We learn

$$\gamma_1 = 2b_{11} = 3, \quad a_1 = -\frac{5}{2}. \quad (23)$$

Next, let us put

$$BF_3 = g \frac{\alpha X^\epsilon}{\epsilon} [B_{31} - 3\epsilon + O(\epsilon^2)] \quad (24)$$

for the moment and consider the orthogonality constraint at $O(g^3)$. Using $BB_0 = 1$, $BF_1 = 0$ we obtain

$$\begin{aligned} \frac{\langle \bar{K}(2) O(1) \rangle_{g^3}}{\langle \bar{B}(2) B(1) \rangle_{g^0}} &= BF_3 - 4gBB_2 - 4g\alpha \frac{b_{11}}{\epsilon} + \frac{gN}{32\pi^2} FF_2 + \frac{gN}{32\pi^2} \alpha \left(\frac{f_{11}}{\epsilon} + f_{10} \right) FF_0 \\ &= g\alpha \left[\frac{1}{\epsilon} (B_{31} + 6 + 4f_{11}) + \log(X) (B_{31} + 12 - 4f_{11}) + (7 - 8f_{11} + 4f_{10}) + O(\epsilon) \right]. \end{aligned} \quad (25)$$

This vanishes up to $O(\epsilon)$ if

$$B_{31} = -9, \quad f_{11} = \frac{3}{4}, \quad f_{10} = -\frac{1}{4}. \quad (26)$$

The singular term f_{11}/ϵ in Z_F is necessary, because we need to solve two different equations in order to eliminate the pole and the logarithm. This fixes both the f_{11} term and the pole term in BF_3 . According to (17) the graph calculation does indeed meet the requirement $B_{31} = -9$. The finite pieces in BF_3 , BB_2 , FF_2 and f_{11} times the subleading order in FF_0 determine the remaining coefficient $f_{10} = -1/4 \neq 0$. Hence in this scheme the Konishi anomaly is affected by renormalisation.

Up to this point we reviewed the analysis of [9]. Next, we look at

$$\begin{aligned} \frac{\langle \bar{K}(2) K(1) \rangle_{g^4}}{\langle \bar{B}(2) B(1) \rangle_{g^0}} &= BB_4 + \alpha \frac{2b_{11}}{\epsilon} BB_2 + \alpha^2 \left(\frac{b_{11}^2}{\epsilon^2} + \frac{2b_{22}}{\epsilon} + \frac{2b_{21}}{\epsilon} \right) + \\ &\quad + 2 \frac{gN}{32\pi^2} BF_3 + \left(\frac{gN}{32\pi^2} \right)^2 FF_2 + 2\alpha \left(\frac{f_{11}}{\epsilon} + f_{10} \right) FF_0 \end{aligned} \quad (27)$$

or by substituting the explicit formulae

$$\begin{aligned} \frac{\langle \bar{K}(2) K(1) \rangle_{g^4}}{\langle \bar{B}(2) B(1) \rangle_{g^0}} &= \alpha^2 \left[\frac{1}{\epsilon^2} \left(B_{42} + 2 b_{22} - \frac{27}{4} \right) + \frac{1}{\epsilon} \log(X) (2 B_{42} - 9) + (\log(X))^2 \left(2 B_{42} - \frac{9}{2} \right) + \right. \\ &\quad \left. + \frac{1}{\epsilon} \left(2 b_{21} + \frac{3}{4} \right) + \log(X) \frac{21}{2} + \left(\frac{17}{2} + \frac{9}{2} \zeta(3) \right) + O(\epsilon) \right] \end{aligned} \quad (28)$$

where we have put

$$BB_4 = \left(\frac{\alpha X^\epsilon}{\epsilon} \right)^2 \left[B_{42} + \frac{45}{4} \epsilon + \left(\frac{45}{4} + \frac{63}{2} \zeta(3) \right) \epsilon^2 + \dots \right] \quad (29)$$

for now. The elimination of the singular terms leads to three conditions, one of which we solve for B_{42} . Upon equating with (22) we find

$$B_{42} = \frac{9}{2}, \quad b_{22} = \frac{9}{8}, \quad \gamma_2 = -3, \quad b_{21} = -\frac{3}{8}, \quad a_2 = \frac{17}{2} + \frac{9}{2} \zeta(3). \quad (30)$$

Once again, the leading coefficient in BB_4 stated in (17) does in fact take the right value.

It remains to analyse the orthogonality constraint at $O(g^5)$. The system is overdetermined as before so that we start with

$$BF_5 = g \left(\frac{\alpha X^\epsilon}{\epsilon} \right)^2 [B_{52} + B_{51} \epsilon + (-3 + 90 \zeta(3)) \epsilon^2 + \dots] \quad (31)$$

(The pole part of FF_4 is already constrained by the protectedness of O^{10}). We put in the other correlators from (17) and the coefficients of the Z-factors already derived.

$$\begin{aligned} \frac{\langle \bar{K}(2) O(1) \rangle_{g^5}}{\langle \bar{B}(2) B(1) \rangle_{g^0}} &= g \alpha^2 \left[\frac{1}{\epsilon^2} (B_{52} + 4 f_{22} - 18) + \frac{1}{\epsilon} \log(X) \left(2 B_{52} - 4 f_{22} - \frac{63}{2} \right) + \right. \\ &\quad + (\log(X))^2 \left(2 B_{52} + 2 f_{22} - \frac{135}{4} \right) + \frac{1}{\epsilon} (B_{51} + 4 f_{21} - 8 f_{22} - 30) + \\ &\quad \left. + \log(X) (2 B_{51} - 4 f_{21} + 8 f_{22} - 75) + (4 f_{20} - 8 f_{21} + 4 f_{22} - 7) + O(\epsilon) \right] \end{aligned} \quad (32)$$

The correlator vanishes up to $O(\epsilon)$ if²

$$B_{52} = \frac{33}{2}, \quad B_{51} = 35, \quad f_{22} = \frac{3}{8}, \quad f_{21} = -\frac{1}{2}, \quad f_{20} = \frac{3}{8}. \quad (33)$$

In conclusion, all three $O(\alpha^2)$ mixing coefficients in Z_F are non-vanishing. Interestingly, $\zeta(3)$ cancels from f_{20} . The fact that the sum of the graphs in $\langle \bar{B}F \rangle_{g^5}$ reproduces B_{52} , B_{51} is extremely non-trivial by looking at the orders $1/\epsilon^3$ and $1/\epsilon^2$ in the explicit results in the equations (55) and (60). The constraints imposed by conformal invariance after renormalisation give an excellent test of the graph calculation.

To make touch with the discussion in [9], we write $F = O + 4gB$ and divide K by $1 + \alpha/2 - \alpha^2/8$ in order to eliminate finite contributions from the shifted renormalisation factor $Z_B + \alpha Z_F/2$ which now multiplies B . The renormalised operator mixture fits the general form

$$\begin{aligned} \tilde{K} &= \tilde{Z}_B B + \frac{gN}{32\pi^2} \tilde{Z}_O O \\ \tilde{Z}_B &= 1 + \alpha \frac{\gamma_1}{2\epsilon} + \alpha^2 \left(\frac{\gamma_1^2}{8\epsilon^2} + \frac{\gamma_2}{4\epsilon} \right) + \dots \\ \tilde{Z}_O &= 1 + \alpha \left(\frac{\gamma_1}{4\epsilon} + \omega_1 \right) + \alpha^2 \left(\frac{\gamma_1^2}{24\epsilon^2} + \frac{\gamma_2 + \omega_1 \gamma_1}{6\epsilon} + \omega_2 \right) + \dots \end{aligned} \quad (34)$$

predicted by (14), with

$$\gamma_1 = 3, \quad \gamma_2 = -3, \quad \omega_1 = -\frac{3}{4}, \quad \omega_2 = \frac{7}{8}. \quad (35)$$

Our effort shows that the renormalisation scheme developed in [9] is operational also at the second order in α , and it yields the previously unknown number ω_2 which could be called the two-loop Konishi anomaly.

²We could have determined b_{22} from here, too.

3 $\langle \bar{B}F \rangle_{g^5}$

The complete set of superdiagrams is displayed in Figure 1 and Figure 6. The diagrams in Figure 1 are special in that they have two cubic non-abelian vertices or one non-abelian four-vertex. They all turn out to be derived topologies³ of the first diagram G_0 so that they can conveniently be summed into one effective numerator $\tilde{G}_0$.

G_0




Figure 1

Suppose that we start by Wick contracting only the matter part of the diagrams, i.e. the operator $\bar{B}(x_7)$ on the cubic chiral vertex at point 6 and the respective matter/YM vertices at point 4 and 5, where present. There are three “matter parts”:




Figure 2

The matter line has an arrow pointing from the antichiral to the chiral end. One way of writing this propagator is

$$\langle \bar{\phi}(i) \phi(j) \rangle = D \cdot D \bar{D} \cdot \bar{D} |_i \Pi_{ij}, \quad D \cdot D = -\frac{1}{4} D^\alpha D_\alpha, \quad \bar{D} \cdot \bar{D} = -\frac{1}{4} \bar{D}_{\dot{\alpha}} \bar{D}^{\dot{\alpha}}. \quad (36)$$

Here $\phi(i) = \phi(x_i, \theta_i, \bar{\theta}_i)$, and

$$\Pi_{ij} = \frac{\delta(\theta_{ij}) \delta(\bar{\theta}_{ij})}{-4\pi^2 x_{ij}^2}, \quad \delta(\theta_{ij}) = \theta_{ij}^\alpha \theta_{ij\alpha}, \quad \delta(\bar{\theta}_{ij}) = \bar{\theta}_{ij\dot{\alpha}} \bar{\theta}_{ij}^{\dot{\alpha}} \quad (37)$$

where $x_{ij}, \theta_{ij}, \bar{\theta}_{ij}$ denote coordinate differences. The Yang-Mills propagator is $\langle V(i) V(j) \rangle = -\Pi_{ij}$.

For this class of graphs we will use “ D -algebra” to derive the numerators. The tools are partial integration, the commutation relation $\{D_\alpha, \bar{D}_{\dot{\alpha}}\} = -2i\partial_{\alpha\dot{\alpha}}$ and the shift rule

$$D_i \Pi_{ij} = -D_j \Pi_{ij}. \quad (38)$$

The idea is to free all but one line of each loop from spinor derivatives. Due to the shift identity and the possibility of reordering via the commutation relations, we may reduce to spacetime derivatives and maximally two chiral and two antichiral spinor derivatives on each Π_{ij} . On the last line in each loop we need to keep only terms with all four derivatives, since

$$\delta(\theta_{ij}) \delta(\bar{\theta}_{ij}) (D \cdot D \bar{D} \cdot \bar{D} \delta(\theta_{ij}) \delta(\bar{\theta}_{ij})) = \delta(\theta_{ij}) \delta(\bar{\theta}_{ij}) \quad (39)$$

³topologies that arise by cancelling a line

while less than four derivatives between two δ -functions vanish.

Let us evaluate matter part M_1 with this technique: We absorb the $\bar{D}.\bar{D}$ of the 7-6 line into the measure at the chiral vertex 6 and partially integrate the $D.D$ away from it onto the 4-6 and the 5-6 lines. We then shift all spinor derivatives to points 4 and 5 respectively. Last, we partially integrate the spinor derivatives away from the 4-6 line, onto $V(4)$ and the 7-4 line, likewise at point 5. The result is

$$\begin{aligned}
\frac{M_1}{-6N^2g^3} = & \text{Tr}[(D.D V(4)) V(5)] \square_{46} \square_{57} + \text{Tr}[(D.D V(4)) (\bar{D}_{\dot{\alpha}} D_{\alpha} V(5))] \square_{46} \frac{i}{2} \partial_{57}^{\dot{\alpha}\alpha} - \\
& \text{Tr}[(D.D V(4)) (\bar{D}.\bar{D} D.D V(5))] \square_{46} + \\
& \text{Tr}[V(4) (D.D V(5))] \square_{56} \square_{47} + \text{Tr}[(\bar{D}_{\dot{\alpha}} D_{\alpha} V(4)) (D.D V(5))] \square_{56} \frac{i}{2} \partial_{47}^{\dot{\alpha}\alpha} - \\
& \text{Tr}[(\bar{D}.\bar{D} D.D V(4)) (D.D V(5))] \square_{56} - \\
& \text{Tr}[(D^{\alpha} V(4)) (D_{\alpha} V(5))] \frac{1}{2} \square_{46} \square_{56} - \\
& \text{Tr}[(D_{\alpha} V(4)) (D.D \bar{D}_{\dot{\alpha}} V(5))] \square_{46} \frac{i}{2} \partial_{56}^{\dot{\alpha}\alpha} + \\
& \text{Tr}[(D.D \bar{D}_{\dot{\alpha}} V(4)) (D_{\alpha} V(5))] \square_{56} \frac{i}{2} \partial_{46}^{\dot{\alpha}\alpha} - \\
& \text{Tr}[(D.D \bar{D}_{\dot{\alpha}} V(4)) (D.D \bar{D}^{\dot{\beta}} V(5))] \frac{1}{2} \partial_{46}^{\dot{\alpha}\alpha} \partial_{56\alpha\dot{\beta}} \quad (40)
\end{aligned}$$

In the last formula ∂_{ij} only acts on the $i-j$ line, at point i . For notational convenience we have omitted $\Pi_{46}\Pi_{47}\Pi_{56}\Pi_{57}\Pi_{67}$ and the integrations. Similarly,

$$\begin{aligned}
\frac{M_1}{6N^2g^3} = & \text{Tr}[V(4) (D.D V(5))] \square_{56} \square_{47} + \text{Tr}[(\bar{D}_{\dot{\alpha}} D_{\alpha} V(4)) (D.D V(5))] \square_{56} \frac{i}{2} \partial_{47}^{\dot{\alpha}\alpha} - \\
& \text{Tr}[(\bar{D}.\bar{D} D.D V(4)) (D.D V(5))] \square_{56} + \\
& \text{Tr}[(D.D V(4)) V(5)] \square_{56} \square_{45} - \text{Tr}[(D^{\alpha} V(4)) (D_{\alpha} V(5))] \frac{1}{2} \square_{56} \square_{45} + \\
& \text{Tr}[(D.D \bar{D}_{\dot{\alpha}} V(4)) (D_{\alpha} V(5))] \square_{56} \frac{i}{2} \partial_{45}^{\dot{\alpha}\alpha} \quad (41)
\end{aligned}$$

Next, $\square_{56}$ leads to the same derived topology in M_1 and M_2 (by shrinking the 5-6 line to a point) so that the “numerators” can simply be added:

$$\begin{aligned}
\frac{M_1 + M_2 + (M_2 : (4 \leftrightarrow 5))}{6N^2g^3} = & \text{Tr}[\bar{\lambda}_{\text{lin } \dot{\alpha}}(4) \bar{\lambda}_{\text{lin}}^{\dot{\beta}}(5)] \frac{1}{2} \partial_{46}^{\dot{\alpha}\alpha} \partial_{56\alpha\dot{\beta}} + (\text{Tr}[(D.D V(4)) V(5)] - \\
& \frac{1}{2} \text{Tr}[(D^{\alpha} V(4)) (D_{\alpha} V(5))] + \text{Tr}[V(4) (D.D V(5))]) \square_{46} \square_{56} \quad (42)
\end{aligned}$$

with the linear part of the antichiral physical fermion in the gauge multiplet $\bar{\lambda}_{\text{lin}}^{\dot{\alpha}} = D.D \bar{D}^{\dot{\alpha}} V$. On the other hand:

$$\frac{M_3}{6N^2g^3} = -D.D \text{Tr}[V(4)V(4)] \square_{46} \quad (43)$$

Due to Greens’ function equation $\square_{56} \Pi_{56} = +\delta^8(z_{56})$ this cancels the $\square_{46}\square_{56}$ terms in (42). In Figure 1 we can therefore work with one effective matter part

$$\tilde{M}_1 = 3N^2g^3 \text{Tr}[\bar{\lambda}_{\text{lin } \dot{\alpha}}(4) \bar{\lambda}_{\text{lin}}^{\dot{\beta}}(5)] \partial_{46}^{\dot{\alpha}\alpha} \partial_{56\alpha\dot{\beta}}. \quad (44)$$

Note that putting $\theta_7 = \bar{\theta}_7 = 0$ also selects the $\theta = \bar{\theta} = 0$ component of $\bar{\lambda}_{\text{lin}}$ at both open ends.

To show consistency with [9] we recompute the leading $O(g^3)$ contribution to $\langle \bar{B}F \rangle$. With

$$F = \text{Tr}[\lambda^{\alpha} \lambda_{\alpha}] \quad \lambda^{\alpha} = -\frac{1}{g} \bar{D}.\bar{D} e^{gV} D^{\alpha} e^{-gV}$$

we find

$$\langle \bar{B}(7) F(1) \rangle_{g^3} = -24g^3N^2(N^2 - 1) \partial_{46}^{\dot{\alpha}\alpha} \partial_{56\alpha\dot{\beta}} \partial_{15}^{\dot{\beta}\beta} \partial_{14\beta\dot{\alpha}}. \quad (45)$$

Once again, for notational convenience we only gave the numerator and omitted the integral itself.

Throughout the paper we employ supersymmetric Fermi-Feynman gauge. The last result would in fact have been immediate in Wess-Zumino (WZ) Fermi-Feynman gauge. However, the D-algebra can be done with the aid of a computer programme so that the non-supersymmetric gauge does not offer any advantage for the g^5 part of the correlator.




Figure 3

The first picture in Figure 3 displays the effective matter part $\tilde{M}_1$ contracted onto $F(1)$, or equivalently the one graph in $\langle \bar{B} F \rangle_{g^3}$ in WZ gauge; here the numbers label the spacetime points. The second picture reproduces the momenta assignment from the discussion of three-loop p-integrals in [12], where this topology is called **BU**. Fourier transform leads to the identification

$$\partial_{14} \rightarrow i p_6, \quad \partial_{15} \rightarrow -i p_7, \quad \partial_{46} \rightarrow -i p_4, \quad \partial_{56} \rightarrow -i p_5. \quad (46)$$

The negative sign of the five matter propagators accounts for an extra minus, so that in momentum space

$$\begin{aligned} \langle \bar{B} F \rangle_{g^3} &= -48 g^3 N^2 (N^2 - 1) [(p_4 \cdot p_5)(p_6 \cdot p_7) - (p_4 \cdot p_7)(p_5 \cdot p_6) + (p_4 \cdot p_6)(p_5 \cdot p_7)] \\ &= -\frac{12 g^3 N^2 (N^2 - 1)}{(4\pi)^6} \left[\frac{1}{\epsilon^2} + \frac{28}{3} \frac{1}{\epsilon} + \frac{166}{3} + O(\epsilon^1) \right] q^2 \left(\frac{q^2}{\mu^2} \right)^{-3\epsilon}. \end{aligned} \quad (47)$$

The second line shows the result of *Mincer* in its $\overline{MS}$ mode.⁴

After this digression let us return to the diagrams of Figure 1. Since combinatorics is linear, the collection of ten supergraphs can be viewed as the effective matter part contracted onto the pure Yang-Mills correlator

$$\tilde{Y}_1 = \langle \bar{\lambda}_{\dot{\alpha} \text{lin}}^a(4) \bar{\lambda}_{\text{lin}}^{\dot{\beta} b}(5) F(1) \rangle_{g^2} \quad (48)$$

where a, b are colour indices. The latter contains the following graphs:




Figure 4

The D-algebra for this sector is cumbersome; as an example we sketch a strategy for the first diagram Y_1 . There are three derivatives on each outer leg and four at each vertex. We resorted to a *Mathematica* script for Wick contraction and superspace algebra. One may choose to shift the derivatives to point 2 and 3, respectively, and then clear the 1-2 and 1-3 lines by partial integration. The derivatives on the 2-3 line are subsequently shifted to one end and reordered. Due to formula (39) only terms with $D \cdot D \bar{D} \cdot \bar{D}$ need to be kept. Next, we regroup the derivatives on the 2-4 and the 3-5 legs. Upon putting the outer theta variables to zero only terms with four spinor derivatives on both the 2-4 and the 3-5 legs survive.

In a similar way,

$$\begin{aligned} \frac{Y_2}{2N g^2 \delta^{ab}} &= \frac{7}{2} \left(\delta_{\dot{\alpha}}^{\dot{\beta}} (\square_1 - \square_{15} - \square_{24}) + (\partial_{15}^{\dot{\beta}\beta} \partial_{24\beta\dot{\alpha}} - \partial_{24}^{\dot{\beta}\beta} \partial_{15\beta\dot{\alpha}}) \right) - 3 \delta_{\dot{\alpha}}^{\dot{\beta}} \square_{24}, \\ \frac{Y_4}{2N g^2 \delta^{ab}} &= \delta_{\dot{\alpha}}^{\dot{\beta}} (\square_1 + \frac{2}{3} \square_{15} + \frac{2}{3} \square_{24}) + (\partial_{24}^{\dot{\beta}\beta} \partial_{25\beta\dot{\alpha}} - \partial_{25}^{\dot{\beta}\beta} \partial_{24\beta\dot{\alpha}}) \end{aligned} \quad (49)$$

where ∂_1 is a total derivative w.r.t. x_1 . Clearly Y_2, Y_3, Y_4 arise from Y_1 by $\square_{13}, \square_{12}, \square_{23}$, respectively. Here the Green's function equation brings in an additional minus sign because $\langle V(i)V(j) \rangle = -\Pi_{ij}$. Upon

⁴The definition of the mass-scale $\tilde{\mu}$ is given before equation (55).

adding these contributions into Y_1 :

$$\begin{aligned} \frac{\tilde{Y}_1}{2Ng^2\delta^{ab}} &= 4\Box_1(\partial_{24}^{\dot{\beta}\beta}\partial_{35\beta\dot{\alpha}} - \partial_{35}^{\dot{\beta}\beta}\partial_{24\beta\dot{\alpha}}) - (\Box_{24} + \Box_{35})(\partial_1^{\dot{\beta}\beta}\partial_{23\beta\dot{\alpha}} - \partial_{23}^{\dot{\beta}\beta}\partial_{1\beta\dot{\alpha}}) \\ &\quad - 4\Box_1(\Box_1 - \Box_{24} - \Box_{35}) - (\Box_{24} - \Box_{35})^2 + (\Box_{12}\Box_{35} + \Box_{13}\Box_{24}) \\ &\quad + (\Box_{12}\Box_{24} + \Box_{13}\Box_{35}) - \frac{2}{3}(\Box_{24} + \Box_{35})\Box_{23} \end{aligned} \quad (50)$$

As another cross-check we derive $\langle \bar{F}(4) F(1) \rangle_{g^2}$ from these formulae:

$$\delta_{\beta}^{\dot{\alpha}} \delta^{ab} \tilde{Y}_1(5 \rightarrow 4) = 2g^2 N(N^2 - 1) [-8\Box_1^2 + 7\Box_1(\Box_{24} + \Box_{34}) + 3(\Box_{12}\Box_{34} + \Box_{13}\Box_{24})] \quad (51)$$

When identifying the points tadpole terms were put to zero. Finally, the mirror image of Y_2 , Y_3 arising from the cubic part of $\bar{F}(4)|_g$ and the tree-like contribution from $\langle \bar{F}(4)|_g F(1)|_g \rangle_{g^2}$ have to be added. The result is

$$\langle \bar{F}(4) F(1) \rangle_{g^2} = 32(N^2 - 1) g^2 N (\Box_{12}\Box_{34} - \frac{1}{2}\Box_1^2) \quad (52)$$

consistent with the results in [9] for the $SU(4)$ component $\tilde{F} = \text{Tr}((\nabla^\alpha \Phi^1)(\nabla_\alpha \Phi^1))$ of F^{10} . According to (63) in the next section, the second term in the last formula is a contact contribution proportional to $\zeta(3)$. The two box operators in the first term break the integrations and we find $-32g^2 N(N^2 - 1) \langle \bar{\phi}(4)\phi(1) \rangle^3$ due to the opposite sign of the matter propagator. In the combination $F - 4gB$ this is cancelled by $16g^2 \langle \bar{B}(4)B(1) \rangle_{g^0}$ as required by protectedness.

By putting formulae (44) and (50) together we obtain the desired effective numerator $\tilde{G}_0$ for the first topology in Figure 1. Graph G_0 is of topology **C** in the list in Figure 5 (the numbers in the figure label the momenta $p_1 \dots p_{10}$).




Figure 5

In Section 6 we comment on the realisation of the Laporta algorithm [13] with which we evaluated such four-loop diagrams with rather general six-derivative numerators. In topology **C** we have chosen the momenta q, p_1, p_2, p_3, p_4 as independent. To calculate $\tilde{G}_0$ we identify

$$\partial_1 \rightarrow iq, \quad \partial_{24} \rightarrow ip_1, \quad \partial_{23} \rightarrow -ip_2, \quad \partial_{46} \rightarrow ip_3, \quad \partial_{56} \rightarrow ip_4. \quad (53)$$

Between these one may form ten dot products. In total there are ten interior lines, so that six of the dot products may be eliminated in favour of the four others and the squares of the dependent momenta. In terms of $p_2 \cdot p_3, p_2 \cdot p_4, p_3 \cdot q, p_4 \cdot q$ and $p_1^2 \dots p_{10}^2, q^2$ the effective numerator has 47 terms. Flip symmetries of the original topology and derived diagrams could be used to simplify.

$$\tilde{G}_0 = 12g^5 N^3(N^2 - 1) [8q^2 p_3 \cdot q p_4 \cdot q - 2(p_2 \cdot p_4 p_3 \cdot q - p_2 \cdot p_3 p_4 \cdot q)(p_1^2 + p_9^2) + (42 \text{ terms})] \quad (54)$$

(The overall negative sign from the odd number of matter propagators has been taken into account.) The omitted terms all have at least two distinct complete squares. Each $p_i^2, i = 1 \dots 10$ cancels a line in the denominator and thus leads to a simplification within the Laporta approach (although p_i^4 is generally not better than p_i^2). The first term in the square brackets is thus the hardest to analyse; an algorithm that solves this problem can also calculate the other pieces.

In the $\overline{MS}$ scheme fractional powers of 4 and π and the Euler constant are absorbed into the mass scale by putting $\bar{\mu}^2 = \mu^2 4\pi e^{-\gamma_E}$. One may further scale away $\zeta(2)$ by the redefinition $\tilde{\mu}^{2\epsilon} = \bar{\mu}^{2\epsilon}(1 - \epsilon^2 \pi^2/12 + O(\epsilon^4))$. In this convention our result for the sum of the diagrams in Figure 1 takes the form

$$\tilde{G}_0 = -\frac{12g^5 N^3(N^2 - 1)}{(4\pi)^8} \left[\frac{19}{72} \frac{1}{\epsilon^3} + \frac{53}{16} \frac{1}{\epsilon^2} + \left(\frac{7957}{288} - \frac{53}{4} \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \right] q^2 \left(\frac{q^2}{\tilde{\mu}^2} \right)^{-4\epsilon}. \quad (55)$$

At the first glimpse not all diagrams in Figure 1 can in fact be drawn into topology **C**. We have shown above that the numerators of the matter parts M_2, M_3 have d'Alembert operators acting onto the 5-6

and 4-6 lines, respectively, common to all terms. These are marked by grey boxes in the graphs of Figure 1. Contracting the corresponding lines yields topologies that can also be derived from **C**.

We have indicated such “topology changing” box operators on the diagrams of Figure 6, too. For instance, graphs G_{31} and G_{32} (the one-loop correction to the chiral vertex in matter part M_1) have originally neither of the four topologies **A**, **B**, **C**, **D**. The grey box operators are positioned such that G_{31} can be obtained from **B** by a factor p_1^2 , and G_{32} from **A** by p_7^2 or equivalently from **D** by p_5^2 .

Figure 6 contains several classes of diagrams that could be summed prior to evaluation like the graphs of Figure 1. The graphs $G_{16} \dots G_{20}$, $G_{21} \dots G_{25}$, $G_{26} \dots G_{30}$ and $G_{31}, G_{32}; G_{33}, G_{34}$ and finally G_{40}, G_{41} are all vertex corrections relating to the matter parts M_1, M_2, M_3 contracted onto $F(1)$. Even in the most complicated graphs G_{16}, G_{19}, G_{20} (type **D**) there is at least one square of an interior momentum in each numerator term, hence the complexity is lower than for $\tilde{G}_0$. There is no calculational disadvantage in individually treating the diagrams, so that we separately list the contribution of every supergraph in order to make the whole calculation more verifiable.

The longest numerators for the **A**, **B** topologies arise from the graphs G_3 and G_{10} in Figure 6 with 134 and 183 numerator terms in our momenta assignments, respectively. These as well as the numerators of G_1, G_6, G_7 contain terms with no squared momentum. Yet, in all such terms at least two of the three non-trivial dot products involve the outer momentum q . The numerator reduction can then still be accomplished with a slightly restricted matrix; terms with one or no q are not needed in the ansatz since q cannot be made to disappear by differentiation w.r.t. loop momenta.

In our way of organising the Laporta algorithm by stepwise elimination of lines (see Section 6), topology **A** is the hardest case because it contains only the triangle between p_4, p_8, p_9 . Using the triangle rule (84) we immediately fall upon two derived topologies that must be further reduced by the Laporta ansatz. In topology **B** we use (84) on the triangle consisting of p_4, p_7, p_8 , which produces a new triangle in every term. The step to the eight-propagator level is therefore more direct. Topologies **C** and **D** both contain two disjoint triangles.

Graph G_4 is in the same way a subgraph of G_3 as the matter part M_2 belongs to M_1 ; likewise, G_{19} and G_{29} go together, and G_{10} and G_{24} . On the other hand, G_{24} naturally belongs to a class of vertex corrections, while G_{10} does not. It is thus not clear how to assemble all the superdiagrams in Figure 6 into well-defined classes; likewise G_{11} and its subcase G_{12} could be attributed to either of the three pairs with one non-abelian vertex.

The “D-algebra” technique is not convenient in diagrams with too many matter lines, because according to (36) every such propagator brings in four spinor derivatives. In the diagrams of Figure 6 we have rather evaluated the Yang-Mills part (in G_1 and similar diagrams the two lines attached to $F(1)$, in G_3 and G_{11} etc. the complete gluon part) by D-algebra and then taken the remaining spinor derivatives to obtain the Grassmann expansion at $\theta_1 = \bar{\theta}_1 = 0$. The alternative form of the matter propagator

$$\langle \bar{\phi}(i) \phi(j) \rangle = e^{-i((\theta_i \partial_{x_i} \bar{\theta}_i) + (\theta_j \partial_{x_j} \bar{\theta}_j) - 2(\theta_j \partial_{x_i} \bar{\theta}_i))} \frac{1}{-4\pi^2 x_{ij}^2} \quad (56)$$

easily lends itself to Grassmann expansion. The complete numerators are found by taking the product with the expansion of the Yang-Mills parts and integrating out the spinor variables.⁵

Finally, from the sum of $\tilde{G}_0$ and $G_1 \dots G_{48}$ of Figure 6 / equation (60) we find

$$\langle \bar{B}(7) F(1) \rangle_{g^5} = \frac{12 g^5 N^3 (N^2 - 1)}{(4\pi)^8} \left[\frac{11}{2} \frac{1}{\epsilon^3} + \frac{899}{12} \frac{1}{\epsilon^2} + \left(\frac{15043}{24} + 30 \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \right] q^2 \left(\frac{q^2}{\tilde{\mu}^2} \right)^{-4\epsilon} \quad (57)$$

in momentum space. The backward Fourier transform⁶

$$\frac{1}{(2\pi)^D} \int d^D x e^{-i q \cdot x} \frac{1}{(q^2)^\alpha} = \frac{1}{4\pi^{\frac{D}{2}}} \frac{\Gamma(\frac{D}{2} - \alpha)}{\Gamma(\alpha)} \frac{1}{(x^2)^{\frac{D}{2} - \alpha}} \quad (58)$$

yields $(\tilde{\mu}_x^{2\epsilon} = (\mu^2 \pi e^{\gamma_E})^\epsilon (1 + \epsilon^2 \pi^2 / 12 + O(\epsilon^4))$ in configuration space)

$$\langle \bar{B}(7) F(1) \rangle_{g^5} = -\frac{g^5 N^3 (N^2 - 1)}{(4\pi^2)^5} \left[\frac{33}{\epsilon^2} + \frac{70}{\epsilon} + (-6 + 180 \zeta(3)) + O(\epsilon^1) \right] \frac{1}{(x_{17}^2)^3} (x_{17}^2 \tilde{\mu}_x^2)^{5\epsilon}. \quad (59)$$

⁵We have computed $\tilde{G}_0$ also by diagram-wise evaluation using this technique.

⁶The forward transform lacks the factor $1/(2\pi)^D$ and has the roles of q, x and the sign in the exponent exchanged.




Figure 6

Diagrams with vanishing colour factors (w.r.t. $SU(N)$ gauge group) or one-loop propagator bubbles were omitted. Graphs G_{22} , G_{23} , G_{27} , G_{28} , $G_{33} \dots G_{48}$ do not need numerator reduction by the Laporta ansatz, though the algorithm is needed here, too, to determine some $\mathbf{T}_1$ configurations (see below) with one or two non-integer exponents.

In (60) we give the momentum space result for each graph including all signs and factors. The ϵ expansions in the table must be multiplied by a global factor $-12 g^5 N^3 (N^2 - 1) / (4\pi)^8 * q^2 (q^2/\tilde{\mu}^2)^{-4\epsilon}$.

$$\begin{aligned}
G_1 &= \frac{1}{4} \frac{1}{\epsilon^3} + \left(\frac{101}{24} + \frac{1}{2} \zeta(3) \right) \frac{1}{\epsilon^2} + \left(\frac{2077}{48} + \frac{1}{120} \pi^4 + \frac{13}{4} \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_2 &= -\frac{1}{8} \frac{1}{\epsilon^3} - \left(\frac{113}{48} - \frac{1}{4} \zeta(3) \right) \frac{1}{\epsilon^2} - \left(\frac{843}{32} - \frac{1}{240} \pi^4 - \frac{13}{8} \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_3 &= \frac{1}{8} \frac{1}{\epsilon^4} + \frac{83}{48} \frac{1}{\epsilon^3} + \frac{483}{32} \frac{1}{\epsilon^2} + \left(\frac{20507}{192} - \frac{35}{3} \zeta(3) + 5 \zeta(5) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_4 &= -\frac{1}{8} \frac{1}{\epsilon^4} - \frac{31}{16} \frac{1}{\epsilon^3} - \frac{1765}{96} \frac{1}{\epsilon^2} - \left(\frac{26881}{192} - \frac{73}{6} \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_5 &= -\frac{2}{3} \frac{1}{\epsilon^3} - \frac{37}{6} \frac{1}{\epsilon^2} - \left(\frac{95}{4} - 3 \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_6 &= \frac{1}{4} \frac{1}{\epsilon^3} + \left(\frac{35}{8} - \frac{1}{2} \zeta(3) \right) \frac{1}{\epsilon^2} + \left(\frac{739}{16} - \frac{1}{120} \pi^4 - \frac{29}{4} \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_7 &= -\frac{2}{3} \frac{1}{\epsilon^3} - \left(\frac{21}{2} - \zeta(3) \right) \frac{1}{\epsilon^2} - \left(\frac{401}{4} - \frac{1}{60} \pi^4 - \frac{21}{2} \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_8 &= -\left(\frac{1}{2} - \frac{3}{2} \zeta(3) \right) \frac{1}{\epsilon^2} - \left(\frac{37}{4} - \frac{1}{40} \pi^4 - \frac{67}{4} \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_9 &= -\frac{1}{8} \frac{1}{\epsilon^3} - \frac{113}{48} \frac{1}{\epsilon^2} - \frac{843}{32} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{10} &= \frac{1}{8} \frac{1}{\epsilon^4} + \frac{47}{24} \frac{1}{\epsilon^3} + \frac{1789}{96} \frac{1}{\epsilon^2} + \left(\frac{13543}{96} - \frac{179}{12} \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{11} &= \frac{3}{4} \frac{1}{\epsilon^3} + \frac{93}{8} \frac{1}{\epsilon^2} + \left(\frac{1765}{16} - 3 \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{12} &= -\frac{3}{4} \frac{1}{\epsilon^3} - \frac{87}{8} \frac{1}{\epsilon^2} - \left(\frac{1551}{16} - 6 \zeta(3) + 5 \zeta(5) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{13} &= -\frac{1}{2} \frac{1}{\epsilon^3} - \left(\frac{35}{4} + \zeta(3) \right) \frac{1}{\epsilon^2} - \left(\frac{739}{8} + \frac{1}{60} \pi^4 + \frac{13}{2} \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{14} &= \frac{1}{8} \frac{1}{\epsilon^3} + \left(\frac{59}{16} - \frac{3}{4} \zeta(3) \right) \frac{1}{\epsilon^2} + \left(\frac{1579}{32} - \frac{1}{80} \pi^4 - \frac{51}{8} \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{15} &= \frac{3}{4} \frac{1}{\epsilon^3} + \frac{53}{2} \frac{1}{\epsilon^2} + \frac{405}{4} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{16} &= -\frac{2}{3} \frac{1}{\epsilon^3} - \frac{37}{3} \frac{1}{\epsilon^2} - \left(\frac{815}{6} - 6 \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{17} &= \frac{5}{8} \frac{1}{\epsilon^3} + \frac{183}{16} \frac{1}{\epsilon^2} + \frac{4007}{32} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{18} &= \frac{5}{8} \frac{1}{\epsilon^3} + \frac{187}{16} \frac{1}{\epsilon^2} + \frac{4115}{32} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{19} &= \frac{1}{2} \frac{1}{\epsilon^3} + \frac{47}{6} \frac{1}{\epsilon^2} + \left(\frac{224}{3} - 3 \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{20} &= -\frac{4}{3} \frac{1}{\epsilon^3} - \frac{68}{3} \frac{1}{\epsilon^2} - 233 \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{21} &= \frac{3}{4} \frac{1}{\epsilon^3} + \frac{299}{24} \frac{1}{\epsilon^2} + \frac{1993}{16} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{22} &= -\frac{1}{4} \frac{1}{\epsilon^3} - \frac{57}{8} \frac{1}{\epsilon^2} - \frac{1453}{16} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{23} &= -\frac{1}{2} \frac{1}{\epsilon^3} - \frac{39}{4} \frac{1}{\epsilon^2} - \frac{871}{8} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{24} &= -\frac{1}{8} \frac{1}{\epsilon^4} - \frac{23}{12} \frac{1}{\epsilon^3} - \frac{1787}{96} \frac{1}{\epsilon^2} - \left(\frac{7013}{48} - \frac{91}{6} \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{25} &= \frac{1}{2} \frac{1}{\epsilon^3} + \frac{145}{12} \frac{1}{\epsilon^2} + \frac{1187}{8} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{26} &= \frac{7}{12} \frac{1}{\epsilon^3} + \frac{269}{24} \frac{1}{\epsilon^2} + \left(\frac{5965}{48} - 6 \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{27} &= -\frac{1}{2} \frac{1}{\epsilon^3} - \frac{37}{4} \frac{1}{\epsilon^2} - \frac{817}{8} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{28} &= -\frac{1}{2} \frac{1}{\epsilon^3} - \frac{39}{4} \frac{1}{\epsilon^2} - \frac{871}{8} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{29} &= -\frac{5}{8} \frac{1}{\epsilon^3} - \frac{19}{2} \frac{1}{\epsilon^2} - \left(\frac{2853}{32} - 3 \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{30} &= \frac{7}{6} \frac{1}{\epsilon^3} + \frac{233}{12} \frac{1}{\epsilon^2} + \frac{4729}{24} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{31} &= \frac{1}{4} \frac{1}{\epsilon^3} + \left(\frac{35}{8} - \zeta(3) \right) \frac{1}{\epsilon^2} + \left(\frac{739}{16} - \frac{1}{60} \pi^4 - \frac{21}{2} \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{32} &= \frac{1}{4} \frac{1}{\epsilon^3} + \frac{101}{24} \frac{1}{\epsilon^2} + \frac{2077}{48} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{33} &= -\frac{1}{6} \frac{1}{\epsilon^3} - \frac{59}{12} \frac{1}{\epsilon^2} - \frac{1579}{24} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{34} &= -\zeta(3) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{35} &= -\frac{7}{6} \frac{1}{\epsilon^3} - \frac{121}{12} \frac{1}{\epsilon^2} - \frac{745}{24} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{36} &= \frac{9}{8} \frac{1}{\epsilon^3} + \frac{159}{16} \frac{1}{\epsilon^2} + \frac{1215}{32} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{37} &= \frac{7}{8} \frac{1}{\epsilon^3} + \frac{121}{16} \frac{1}{\epsilon^2} + \frac{745}{32} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{38} &= \frac{1}{8} \frac{1}{\epsilon^3} + \frac{59}{16} \frac{1}{\epsilon^2} + \frac{1579}{32} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{39} &= -16 \zeta(3) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{40} &= -8 \zeta(3) \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{41} &= -\frac{4}{3} \frac{1}{\epsilon^3} - \frac{52}{3} \frac{1}{\epsilon^2} - \frac{412}{3} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{42} &= \frac{4}{3} \frac{1}{\epsilon^3} + \frac{52}{3} \frac{1}{\epsilon^2} + \frac{412}{3} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{43} &= -\frac{16}{3} \frac{1}{\epsilon^3} - 64 \frac{1}{\epsilon^2} - \frac{1408}{3} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{44} &= \frac{1}{6} \frac{1}{\epsilon^2} + \frac{13}{4} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{45} &= -\frac{5}{4} \frac{1}{\epsilon^3} - \frac{111}{8} \frac{1}{\epsilon^2} - \frac{1391}{16} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{46} &= \frac{10}{9} \frac{1}{\epsilon^3} + \frac{130}{9} \frac{1}{\epsilon^2} + \frac{1030}{9} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{47} &= -\frac{10}{9} \frac{1}{\epsilon^3} - \frac{130}{9} \frac{1}{\epsilon^2} - \frac{1030}{9} \frac{1}{\epsilon} + O(\epsilon^0) \\
G_{48} &= -\frac{20}{9} \frac{1}{\epsilon^3} - \frac{80}{3} \frac{1}{\epsilon^2} - \frac{1760}{9} \frac{1}{\epsilon} + O(\epsilon^0)
\end{aligned} \tag{60}$$

4 $\langle \bar{B}B \rangle_{g^4}$

At tree-level

$$\langle \bar{B}(2) B(1) \rangle_{g^0} = -\frac{2N(N^2 - 1)}{(4\pi^2)^3} \frac{1}{(x_{12}^2)^3} (x_{12}^2 \tilde{\mu}_x^2)^{3\epsilon} (1 + O(\epsilon^3)) \quad (61)$$

where the $\epsilon^3 \zeta(3) + \dots$ correction arises from $\Gamma(1 - \epsilon)$ in the numerator of the modified propagator in dimensional regularisation. At order g^2 there are two graphs:

$$\langle \bar{B}(2) B(1) \rangle_{g^2} = 12 g^2 N^2 (N^2 - 1) \langle \bar{\phi}(2) \phi(1) \rangle_{g^0} (H_1 + H_2) \quad (62)$$




Figure 7

$$\begin{aligned} H_1 &= -\frac{1}{(4\pi)^4} \left[\frac{1}{\epsilon^2} + \frac{4}{\epsilon} + 12 + O(\epsilon) \right] \left(\frac{q^2}{\tilde{\mu}^2} \right)^{-2\epsilon}, \\ H_2 &= -\frac{1}{(4\pi)^4} [3\zeta(3) + O(\epsilon)] \left(\frac{q^2}{\tilde{\mu}^2} \right)^{-2\epsilon} \end{aligned} \quad (63)$$

whereby in configuration space

$$\langle \bar{B}(2) B(1) \rangle_{g^2} = -\frac{g^2 N}{4\pi^2} \langle \bar{B}(2) B(1) \rangle_{g^0} \left[\frac{3}{\epsilon} + 3 + 9\zeta(3)\epsilon + O(\epsilon^2) \right] (x_{12}^2 \tilde{\mu}_x^2)^\epsilon. \quad (64)$$

The diagram H_2 is of topology $\mathbf{T}_1$ in the nomenclature of [12]. Its numerator is just an outer box operator so that we can take over the result (87) elaborated below. The leading term is of order ϵ^0 in momentum space, which means $O(\epsilon)$ in configuration space. It is therefore a contact term [2]. Note that the contact contribution in (52) is exactly H_2 with a second outer box operator, which does not touch upon the order in ϵ of the leading term.

On the other hand, w.r.t. the $O(g^4)$ part of $\langle \bar{B}B \rangle$ we are only interested in the singular part in momentum space, or equivalently the singular and finite pieces in configuration space. On comparing to a certain protected correlator we can spare most of the work: Expanding in N , protected correlation functions contain several linear combinations of graphs in which the x space singular and finite terms must cancel. One such sum of graphs is also present in $\langle \bar{B}B \rangle_{g^4}$, with identical relative coefficients for the relevant graphs. Using the work of [14]:

$$\frac{\langle \bar{\phi}(2) \phi(1) \rangle_{g^0}}{12 g^4 N^3 (N^2 - 1)} \langle \bar{B}(2) B(1) \rangle_{g^4} = \langle \bar{\phi}(2) \phi(1) \rangle_{g^0} \Sigma_1 + H_2^2 - \Sigma_2 \quad (65)$$

where Σ_1 contains the six four-loop graphs $H_3 \dots H_8$




Figure 8

and

$$\Sigma_2 = \left[\frac{\langle \bar{P}(2) P(1) \rangle_{g^4}}{12 g^4 (N^2 - 1)} \right]_{N^4}, \quad P = \text{Tr}(\phi^1 \phi^1 \phi^2 \phi^2) + \frac{1}{2} \text{Tr}(\phi^1 \phi^2 \phi^1 \phi^2) \quad (66)$$

is a sum of graphs in which only a contact contribution survives because P is an $SU(4)$ component of the half BPS operator $\text{Tr}(\phi^1 \phi^1 \phi^1 \phi^1)$. Hence in x space

$$\langle \bar{B}(2) B(1) \rangle_{g^4} = 12 g^4 N^3 (N^2 - 1) \Sigma_1 + O(\epsilon). \quad (67)$$

The first five graphs in Σ_1 are simple to evaluate with the methods developed for $\langle \bar{B}F \rangle_{g^5}$. For convenience we used *Mincer* for the three-loop ladder subgraph in H_8 . Including all signs and factors

$$\begin{aligned}
H_3 &= \frac{1}{\epsilon^3} + 14 \frac{1}{\epsilon^2} + 121 \frac{1}{\epsilon} + O(\epsilon^0) \\
H_4 &= \frac{1}{4} \frac{1}{\epsilon^3} + \frac{33}{8} \frac{1}{\epsilon^2} + \left(\frac{665}{16} + 6 \zeta(3) \right) \frac{1}{\epsilon} + O(\epsilon^0) \\
H_5 &= -\frac{1}{4} \frac{1}{\epsilon^3} - \frac{33}{8} \frac{1}{\epsilon^2} - \frac{657}{16} \frac{1}{\epsilon} + O(\epsilon^0) \\
H_6 &= \frac{9}{2} \zeta(3) \frac{1}{\epsilon} + O(\epsilon^0) \\
H_7 &= \frac{3}{4} \frac{1}{\epsilon^3} + \frac{87}{8} \frac{1}{\epsilon^2} + \frac{1551}{16} \frac{1}{\epsilon} + O(\epsilon^0) \\
H_8 &= -\frac{1}{4} \frac{1}{\epsilon^3} - \frac{31}{8} \frac{1}{\epsilon^2} - \frac{591}{16} \frac{1}{\epsilon} + O(\epsilon^0)
\end{aligned} \tag{68}$$

to be multiplied by an overall $12 g^4 N^3 (N^2 - 1) / (4\pi)^8 * q^2 (q^2 / \tilde{\mu}^2)^{-4\epsilon}$. Summing up and translating to x space

$$\langle \bar{B}(2) B(1) \rangle_{g^4} = \left(\frac{g^2 N}{4\pi^2} \right)^2 \langle \bar{B}(2) B(1) \rangle_{g^0} \left[\frac{9}{2} \frac{1}{\epsilon^2} + \frac{45}{4} \frac{1}{\epsilon} + \left(\frac{45}{4} + \frac{63}{2} \zeta(3) \right) + O(\epsilon) \right] (x_{12}^2 \tilde{\mu}_x^2)^{2\epsilon}. \tag{69}$$

5 $\langle \bar{\mathcal{O}} \mathcal{O} \rangle_{g^4}$ and $\langle \bar{F} F \rangle_{g^4}$

Let us first consider the two-point function of the half BPS operator

$$\mathcal{O} = \text{Tr}(\phi^1 \phi^1).$$

The two leading orders are

$$\langle \bar{\mathcal{O}}(2) \mathcal{O}(1) \rangle = 2(N^2 - 1) [\langle \bar{\phi}(2) \phi(1) \rangle_{g^0}^2 + 4 g^2 N H_2 + O(g^4)] \tag{70}$$

while the $O(g^4)$ part receives contributions by the following graphs:




Figure 9

The blob in I_5 means the one-loop correction to the cubic vertex $g \int \text{Tr}(\Phi^I [V, \bar{\Phi}_I])$ and the blob in I_6 denotes the two-loop correction to the matter propagator.

$$\begin{aligned}
I_1 &= \zeta(3) \frac{1}{\epsilon} + \frac{\pi^4}{60} + 6 \zeta(3) + O(\epsilon) \\
I_2 &= \zeta(3) \frac{1}{\epsilon} + \frac{\pi^4}{60} + 6 \zeta(3) + 5 \zeta(5) + O(\epsilon) \\
I_3 &= 20 \zeta(5) + O(\epsilon) \\
I_4 &= -30 \zeta(5) + O(\epsilon) \\
I_5 &= -4 \zeta(3) \frac{1}{\epsilon} - \frac{\pi^4}{15} - 24 \zeta(3) + 30 \zeta(5) + O(\epsilon) \\
I_6 &= 2 \zeta(3) \frac{1}{\epsilon} + \frac{\pi^4}{30} + 12 \zeta(3) + O(\epsilon)
\end{aligned} \tag{71}$$

The ϵ expansions in the table come with an overall factor $8 g^4 N^2 (N^2 - 1) / (4\pi)^6 * (q^2 / \tilde{\mu}^2)^{-3\epsilon}$. The sum of the lines in (71) is $25 \zeta(5) + O(\epsilon)$; hence only a contact term remains as required by protectedness.

We only give the sum of the propagator and vertex corrections because the loop order here is not higher than three so that we could use the *Mincer* programme to compute the graphs. Moreover, the vanishing of the simple pole in the sum has been checked in [2].

Note that $I_1 \dots I_3$ are only of order ϵ^{-1} ; in x space they are therefore individually finite, while I_4 is by itself a contact term. Likewise, the five graphs contributing to I_5 are all separately finite in x space. In $\mathcal{N} = 4$ SYM non-protected operators have to be renormalised only due to diagrams with subdivergences coming from the outer ends; as typical examples we mention the pictures of H_1 or H_3 , H_5 above. For protected operators these divergences are absent. Due to the improved power counting in superspace,

connected graphs tend to be finite when they have sufficiently many outer legs. On the other hand, the individual graphs contributing to the (overall finite) two-loop matter blob have poles up to second order in momentum space which the “kinematical” third loop in I_6 even promotes to $1/\epsilon^3$.

Collecting terms, in configuration space

$$\begin{aligned} \langle \bar{\mathcal{O}}(2) \mathcal{O}(1) \rangle = \langle \bar{\mathcal{O}}(2) \mathcal{O}(1) \rangle_{g^0} & \left[1 - \alpha \left(6 \zeta(3) \epsilon + O(\epsilon^2) \right) (x_{12}^2 \tilde{\mu}_x^2)^\epsilon \right. \\ & \left. + \alpha^2 \left(\frac{75}{4} \zeta(5) \epsilon + O(\epsilon^2) \right) (x_{12}^2 \tilde{\mu}_x^2)^{2\epsilon} + O(\alpha^3) \right], \quad \alpha = \frac{g^2 N}{4\pi^2}. \end{aligned} \quad (72)$$

Next, writing $\mathcal{O} = \text{Tr}[(e^{-gV} \phi^1 e^{gV}) (e^{-gV} \phi^1 e^{gV})]$ and using the equation of motion

$$D.D (e^{-gV} \phi^1 e^{gV}) = g [\bar{\phi}_2, \bar{\phi}_3] \quad (73)$$

we derive

$$\begin{aligned} \frac{1}{2} D^\alpha D_\alpha \mathcal{O} &= \check{F} - 4g \check{B}, \\ \check{F} &= \text{Tr}[(D^\alpha (e^{-gV} \phi^1 e^{gV})) (D_\alpha (e^{-gV} \phi^1 e^{gV}))], \\ \check{B} &= \text{Tr}[(e^{-gV} \phi^1 e^{gV}) [\bar{\phi}_2, \bar{\phi}_3]]. \end{aligned}$$

Recall that the F involving two Yang-Mills fermions which we used before and the representative $\check{F}$ are different $SU(4)$ components of the same operator, likewise for $\check{B}$ and its $\mathcal{N} = 1$ chiral companion $B = \text{Tr}(\phi^1 [\phi^2, \phi^3])$.

The Grassmann expansion of the correlator $\langle \bar{\mathcal{O}}(2) \mathcal{O}(1) \rangle$ is given by the same exponential shift operator as for the superspace matter propagator:

$$\langle \bar{\mathcal{O}}(2) \mathcal{O}(1) \rangle = e^{i((\theta_1 \partial_1 \bar{\theta}_1) + (\theta_2 \partial_1 \bar{\theta}_2) - 2(\theta_1 \partial_1 \bar{\theta}_2))} \langle \bar{\mathcal{O}}(2) \mathcal{O}(1) \rangle_{\theta_{1,2} = \bar{\theta}_{1,2} = 0} \quad (74)$$

It follows that

$$\langle (\check{F} - 4g \check{B})(1) (\check{F} - 4g \check{B})(2) \rangle = 4 D.D|_1 \bar{D}.\bar{D}|_2 \langle \bar{\mathcal{O}}(2) \mathcal{O}(1) \rangle = -4 \square_1 \langle \bar{\mathcal{O}}(2) \mathcal{O}(1) \rangle. \quad (75)$$

In particular, for the $\theta = \bar{\theta} = 0$ component we should find to leading order in ϵ

$$\begin{aligned} & \langle (\check{F} - 4g \check{B})(1) (\check{F} - 4g \check{B})(2) \rangle \\ &= -\frac{64(N^2 - 1)}{(4\pi^2)^2 (x_{12}^2)^3} (x_{12}^2 \tilde{\mu}_x^2)^{2\epsilon} \left[(1 + O(\epsilon)) - \alpha \epsilon 6 \zeta(3) (x_{12}^2 \tilde{\mu}_x^2)^\epsilon + \alpha^2 \epsilon \frac{75}{4} \zeta(5) (x_{12}^2 \tilde{\mu}_x^2)^{2\epsilon} + \dots \right]. \end{aligned} \quad (76)$$

The tree-level part is obviously right for both F and $\check{F}$, whereas the $O(\alpha)$ contact term agrees with (52) because F and $\check{F}$ have interchangeable two-point functions by R-symmetry invariance. We now wish to verify the $O(\alpha^2)$ contribution in the last equation by a direct graph calculation. We prefer $\check{F}$ for this purpose because $F = \lambda^\alpha \lambda_\alpha + \dots$ would, of course, heavily involve the Yang-Mills sector, while we have seen above that the D-algebra creates quite some work already at $O(\alpha)$.

Upon expanding the exponentials in the definition in (4)

$$\begin{aligned} \check{F} &= \text{Tr} \left(D^\alpha \phi^1 D_\alpha \phi^1 - 2g [D^\alpha V, \phi^1] D_\alpha \phi^1 + \right. \\ & \quad \left. g^2 [D^\alpha V, \phi^1] [D_\alpha V, \phi^1] + g^2 [[D^\alpha V, V], \phi^1] D_\alpha \phi^1 + O(g^3) \right). \end{aligned} \quad (77)$$

In the last formula D^α acts only on the field immediately after it. In other words, in the given frame the chiral spinor derivative becomes Yang-Mills covariantized:

$$\nabla^\alpha = D^\alpha - g [D^\alpha V, \bullet] + \frac{g^2}{2} [[D^\alpha V, V], \bullet] + O(g^3) \quad (78)$$

(On the other hand, the constraint $\bar{D} \phi^1 = 0$ is not modified.)

The top graphs in $\langle \check{F} \check{F} \rangle_{g^4, \theta = \bar{\theta} = 0}$ are like in Figure 9 but with a partial spinor derivative on each outer leg. These graphs have the ϵ expansions $\tilde{I}_1 \dots \tilde{I}_6$ in the table below, which are visibly not overall

derivatives of $I_1 \dots I_6$. Some new subdivergences have to be compensated by diagrams arising from the higher terms in (77). Putting $\theta = \bar{\theta} = 0$ at the outer points removes supergraphs with more than two outer V fields. Further, most ways of placing $D, \bar{D}$ on the fields at the outer points lead to vanishing results. The remaining extra diagrams are given in Figure 10. We have put the derivatives onto the graphs where their position could be ambiguous.




Figure 10

$$\begin{aligned}
\tilde{I}_1 &= -\frac{1}{6} \frac{1}{\epsilon^2} - \left(\frac{4}{3} - \frac{1}{2}\zeta(3)\right) \frac{1}{\epsilon} - \frac{19}{3} + \frac{\pi^4}{120} + 3\zeta(3) + O(\epsilon) \\
\tilde{I}_2 &= -\frac{1}{6} \frac{1}{\epsilon^2} - \left(2 - \frac{3}{2}\zeta(3)\right) \frac{1}{\epsilon} - 15 + \frac{\pi^4}{40} + 7\zeta(3) + \frac{5}{2}\zeta(5) + O(\epsilon) \\
\tilde{I}_3 &= \frac{2}{3} \frac{1}{\epsilon^3} + \frac{16}{3} \frac{1}{\epsilon^2} + \left(\frac{88}{3} - 12\zeta(3)\right) \frac{1}{\epsilon} + \frac{416}{3} - \frac{\pi^4}{5} - \frac{196}{3}\zeta(3) + 10\zeta(5) + O(\epsilon) \\
\tilde{I}_4 &= \frac{1}{2} \frac{1}{\epsilon^3} + 5 \frac{1}{\epsilon^2} + \frac{131}{4} \frac{1}{\epsilon} + \frac{711}{4} - \frac{41}{2}\zeta(3) - 15\zeta(5) + O(\epsilon) \\
\tilde{I}_5 &= -\frac{1}{3} \frac{1}{\epsilon^3} - \frac{10}{3} \frac{1}{\epsilon^2} - \left(\frac{87}{4} + 2\zeta(3)\right) \frac{1}{\epsilon} - \frac{1409}{12} - \frac{\pi^4}{30} - \frac{10}{3}\zeta(3) + 15\zeta(5) + O(\epsilon) \\
\tilde{I}_6 &= +\zeta(3) \frac{1}{\epsilon} + \frac{\pi^4}{60} + 7\zeta(3) + O(\epsilon) \\
\tilde{I}_7 &= -2 \frac{1}{\epsilon^3} - 18 \frac{1}{\epsilon^2} - \left(\frac{328}{3} - 12\zeta(3)\right) \frac{1}{\epsilon} - \frac{1684}{3} + \frac{\pi^4}{5} + 110\zeta(3) + O(\epsilon) \\
\tilde{I}_8 &= \frac{1}{6} \frac{1}{\epsilon^2} + (2 - \zeta(3)) \frac{1}{\epsilon} + 15 - \frac{\pi^4}{60} - 3\zeta(3) + O(\epsilon) \\
\tilde{I}_9 &= -\frac{1}{6} \frac{1}{\epsilon^3} - \frac{3}{2} \frac{1}{\epsilon^2} - 9 \frac{1}{\epsilon} - \frac{136}{3} + \frac{17}{6}\zeta(3) + O(\epsilon) \\
\tilde{I}_{10} &= -\frac{1}{4} \frac{1}{\epsilon^3} - \frac{5}{2} \frac{1}{\epsilon^2} - \left(\frac{49}{3} + \zeta(3)\right) \frac{1}{\epsilon} - \frac{265}{3} - \frac{\pi^4}{60} + \frac{13}{4}\zeta(3) + O(\epsilon) \\
\tilde{I}_{11} &= \frac{1}{6} \frac{1}{\epsilon^3} + \frac{5}{3} \frac{1}{\epsilon^2} + \left(11 + \frac{1}{2}\zeta(3)\right) \frac{1}{\epsilon} + \frac{181}{3} + \frac{\pi^4}{120} - \frac{4}{3}\zeta(3) + O(\epsilon) \\
\tilde{I}_{12} &= \frac{1}{3} \frac{1}{\epsilon^3} + 3 \frac{1}{\epsilon^2} + \frac{55}{3} \frac{1}{\epsilon} + 95 - \frac{29}{3}\zeta(3) + O(\epsilon) \\
\tilde{I}_{13} &= \frac{1}{3} \frac{1}{\epsilon^3} + 3 \frac{1}{\epsilon^2} + 18 \frac{1}{\epsilon} + \frac{272}{3} - \frac{23}{3}\zeta(3) + O(\epsilon) \\
\tilde{I}_{14} &= \frac{1}{6} \frac{1}{\epsilon} + \frac{13}{6} - \zeta(3) + O(\epsilon) \\
\tilde{I}_{15} &= \frac{1}{6} \frac{1}{\epsilon} + \frac{13}{6} + O(\epsilon) \\
\tilde{I}_{16} &= \frac{1}{2}\zeta(3) \frac{1}{\epsilon} + \frac{\pi^4}{120} + \frac{7}{2}\zeta(3) + O(\epsilon) \\
\tilde{I}_{17} &= \frac{1}{2} \frac{1}{\epsilon^3} + \frac{9}{2} \frac{1}{\epsilon^2} + \frac{55}{2} \frac{1}{\epsilon} + \frac{285}{2} - \frac{29}{2}\zeta(3) + O(\epsilon) \\
\tilde{I}_{18} &= \frac{1}{12} \frac{1}{\epsilon^3} + \frac{5}{6} \frac{1}{\epsilon^2} + \frac{11}{2} \frac{1}{\epsilon} + \frac{181}{6} - \frac{29}{12}\zeta(3) + O(\epsilon) \\
\tilde{I}_{19} &= \frac{1}{6} \frac{1}{\epsilon^3} + \frac{5}{3} \frac{1}{\epsilon^2} + 11 \frac{1}{\epsilon} + \frac{181}{3} - \frac{29}{6}\zeta(3) + O(\epsilon)
\end{aligned} \tag{79}$$

In the table we have omitted the overall factor $F_0 = 64 g^4 N^2 (N^2 - 1) / (4\pi)^6 * q^2 (q^2 / \tilde{\mu}^2)^{-3\epsilon}$. The sum of $\tilde{I}_1 \dots \tilde{I}_{19}$ yields the first line in the next equation.

$$\begin{aligned}
\langle \bar{F} F \rangle_{g^4} &= F_0 \left[-\frac{1}{2} \frac{1}{\epsilon^2} - \frac{4}{\epsilon} - 19 + 3\zeta(3) + \frac{25}{2}\zeta(5) + O(\epsilon) \right], \\
-8g \langle \bar{F} B \rangle_{g^3} &= F_0 \left[+\frac{3}{2} \frac{1}{\epsilon^2} + \frac{14}{\epsilon} + 83 + O(\epsilon) \right], \\
16g^2 \langle \bar{B} B \rangle_{g^2} &= F_0 \left[-\frac{1}{\epsilon^2} - \frac{10}{\epsilon} - 64 - 3\zeta(3) + O(\epsilon) \right].
\end{aligned} \tag{80}$$

Upon adding up

$$\langle (\bar{F} - 4g\bar{B})(F - 4gB) \rangle_{g^4} = F_0 \frac{25}{2} \zeta(5) + O(\epsilon) \quad (81)$$

which reproduces the $O(\alpha^2)$ term in (76) after Fourier transform.

In conclusion, at the given orders the operator $F - 4gB$ does not need renormalisation. Its anomalous dimension is zero at one and two loops. At loop level the two-point function does pick up contact contributions; remarkably the $O(\alpha)$ contact term has normalisation proportional to $\zeta(3)$ while the $O(\alpha^2)$ term comes with $\zeta(5)$. The equation of motion relating the operator $O = \text{Tr}(\phi^1 \phi^1)$ to $F - 4gB$ is apparently purely classical: To leading order in ϵ the normalisation of the contact part is compatible with superspace differentiation.

6 Integration by parts (IBP) in dimensional regularisation

We follow the last reference in [12] in exposing the fundamental idea. Suppose that any given diagram has a triangle subgraph like the first picture in Figure 11.




Figure 11

The integral associated to the subdiagram is

$$I(\alpha_0, \beta_1, \beta_2, \alpha_1, \alpha_2) = \int d^D p \frac{1}{(p^2)^{\alpha_0} ((p+p_1)^2)^{\beta_1} (p_1^2)^{\alpha_1} ((p+p_2)^2)^{\beta_2} (p_2^2)^{\alpha_2}}. \quad (82)$$

Discarding boundary terms

$$\int d^D p \partial_{p_\mu} \frac{p_\mu}{(p^2)^{\alpha_0} ((p+p_1)^2)^{\beta_1} (p_1^2)^{\alpha_1} ((p+p_2)^2)^{\beta_2} (p_2^2)^{\alpha_2}} = 0 \quad (83)$$

because the integrand is a total derivative. By working out the differentiation:

$$\begin{aligned} I(\alpha_0, \beta_1, \beta_2, \alpha_1, \alpha_2) (D - 2\alpha_0 - \beta_1 - \beta_2) = \\ \beta_1 (I(\alpha_0 - 1, \beta_1 + 1, \beta_2, \alpha_1, \alpha_2) - I(\alpha_0, \beta_1 + 1, \beta_2, \alpha_1 - 1, \alpha_2)) + \\ \beta_2 (I(\alpha_0 - 1, \beta_1, \beta_2 + 1, \alpha_1, \alpha_2) - I(\alpha_0, \beta_1, \beta_2 + 1, \alpha_1, \alpha_2 - 1)) \end{aligned} \quad (84)$$

The generalisation of the formula for polynomial numerators $p^{\nu_1} \dots p^{\nu_n}$ is straightforward. In every term on the r.h.s. of (84) the exponent of one of the propagators of the triangle's top line is diminished by one. To illustrate the use of the formula we compute the integral $\mathbf{T}_1$ (without numerator) given in the second picture of Figure 11. Using (84) on the left triangle in the diagram produces an overall factor $1/\epsilon$ times the third minus the fourth picture. Our convention is that a line without any extra symbol has exponent equal to one. The triangle rule can remove such a line, but in exchange it augments the weight of one of the β lines by one. This is customarily denoted by a dot. The resulting new diagrams are both elementary: One is a convolution, the other the product of two one-loop integrals. The elementary one-loop building block is

$$\int \frac{d^D p}{(2\pi)^D} \frac{1}{(p^2)^\alpha ((q-p)^2)^\beta} = \frac{1}{(4\pi)^2} G(\alpha, \beta) (q^2)^{D/2-\alpha-\beta}, \quad D = 4 - 2\epsilon \quad (85)$$

where

$$G(\alpha, \beta) = (4\pi)^\epsilon R(\alpha) R(\beta) R(D - \alpha - \beta), \quad R(\alpha) = \frac{\Gamma(D/2 - \alpha)}{\Gamma(\alpha)}. \quad (86)$$

For the generalisation to integrals with numerator polynomials we refer to [12]. It follows

$$\mathbf{T}_1 = \frac{1}{\epsilon} G(1, 1) (G(2, 1 + \epsilon) - G(2, 1)) = \frac{1}{(4\pi)^4} \left(6\zeta(3) + \left(\frac{\pi^4}{10} + 12\zeta(3) \right) \epsilon + \dots \right) \frac{1}{q^2} \left(\frac{q^2}{\tilde{\mu}^2} \right)^{-2\epsilon} \quad (87)$$

in the $\overline{MS}$ convention explained above. Almost all three-loop “p-integrals” (propagator type) can be calculated by this trick [12]: By way of example, to solve the **BU** topology displayed in Figure 3 one may start with one of the visible triangles and then iterate the procedure on the resulting **T**₁ (sub)diagrams.

Let us now turn to the four-loop topology **A** of our $\langle \bar{B}F \rangle_{g^5}$ problem. We can use the rule of the triangle once:




Figure 12

The two new topologies contain no further triangle. Nonetheless, the IBP technique can be pushed further by the Laporta ansatz [13]: The integral in the middle of Figure 12 now has propagators with power one for, say, p_1 to p_8 and of power two for p_9 . The most complicated numerators we encounter have three non-trivial dot products. We write the complete set of identities

$$\int d^D p_1 \dots d^D p_4 \partial_{p_i^\mu} \frac{p_j^\mu \text{num}}{p_1^2 \dots p_8^2 (p_9^2)^2} = 0 \quad (88)$$

where $i \in \{1 \dots 4\}$ and p_j can be one of the loop momenta or the outer momentum q . Between the four loop momenta and q one can form fifteen independent square invariants. Let the first ten be the squares of all the interior momenta and of q , then we have to choose five further mixed dot products. Mixed products containing q bring an advantage. We write the 20 identities (88) for any such numerator *num* built from three square invariants that does not by itself cancel a line in the denominator (thus it can contain the mixed dot products, q^2 or one power of p_9^2). Next the differentiation is worked out just as in the case of the triangle rule. The result is a large homogeneous linear system for a basis of integrals, which one may reduce by Gaussian elimination. The matrix is initially very sparse and the elimination has the surprising property that the relative order in ϵ between the terms in the same line remains relatively stable throughout the steps of the algorithm.

Due to memory limitations (1 GB on a power PC and later a Xeon, of which we needed only about one half) we organised the task in a recursive way: In the first step, except for the integral with a sought numerator, all nine-propagator structures are eliminated. We obtain a linear equation relating it to cases with eight or less propagators (the cancellation of a line works as in the case of the triangle rule). The set of eight-propagator configurations is given in Figure 13. The differentiation in (88) produces a second dot, which we did not indicate on the pictures because it can be placed on any of the interior lines. By dimensionality the eight-propagator integrals still have numerators with maximally three kinematic invariants, or less if there is a further cancellation with a denominator term.




Figure 13

On this level, M_{35} and M_{36} once again need numerator reduction by the Laporta ansatz. There is a new feature arising here: Apart from the desired numerator term one can eliminate all eight-propagator cases but one. The conventional choice for the remaining integral is M_{35} or M_{36} , respectively, with trivial numerator q^2 and all exponents equal to one. These integrals are called “masters”. A priori they have to be calculated by independent means. The attempt to eliminate a master usually trivializes the system of equations.

The other integrals in Figure 13 and the seven-propagator configurations found by cancelling a line from M_{35} , M_{36} either have triangles or they are trivial like the last picture in Figure 13. However,

according to equation (85) a one-loop subintegral leaves behind a propagator with non-integer dimension; in Figure 14 we marked this by a cross. If this affects one of the α lines of a triangle subgraph, equation (84) ceases to be helpful. As a consequence, in the last step a variety of $\mathbf{T}_1$ cases with modified propagators again have to be attacked by the Laporta idea. One can eventually backsubstitute starting from a general result for $\mathbf{T}_1(1 + a_1 \epsilon, 1 + a_2 \epsilon, 1 + a_3 \epsilon, 1 + a_4 \epsilon, 1 + a_5 \epsilon)$ due to [15] and explicit results for the masters M_{35}, M_{36} [16].

The integral in the third picture in Figure 12 can be dealt with in the same manner: The bubble integral leads to the three-loop **NO** topology with a cross on one of the outer lines. Once again, the Laporta idea is needed to further reduce this. Foreseeably, there is a master integral for which we choose $\text{NO}(\epsilon, 1, 1, 1, 1, 1, 1)$. In Figure 14 we list the more trivial seven-propagator configurations, of which only the first case needs Laporta reduction.




Figure 14

Finally, topologies **B**, **C**, **D** are somewhat nicer in that one can always get to the eight-propagator level by two successive applications of the triangle rule. They contain the same master integrals M_{35}, M_{36} and only a **BU** topology with a cross on the p_4 or p_5 propagator (c.f. Figure 3) comes in additionally.

The recursive procedure is quite laborious because a separate routine is needed for the evaluation of every triangle solvable derived topology with a numerator. Further, we had to create tables of M_{35} and M_{36} with two dots and three dot products in the numerator, likewise for the other non-elementary cases. These M_{35}, M_{36} integrals present a formidable computing challenge because the matrix size is of the order $1650 * 4400$ in polynomials in ϵ . It seems more appealing to attempt a complete numerator reduction directly on the topologies **A**, **B**, **C**, **D** because there are no dots and one needs to consider only four distinct mixed products in the numerator so that the matrix results considerably smaller.

The advantage of our architecture is that the Gaussian elimination need not be pushed to the end; it was usually sufficient to eliminate less than half of the integrals in a basis to get definite results for any given numerator term. We realised the Gauss algorithm between *ANSI C* and *Mathematica*. The whole problem addressed in this article could be solved on desktop machines with moderate memory, although the total runtime amounted to several weeks. A sample calculation with the programmes will be presented in a separate publication, probably on the example of M_{36} .

We conclude this section with a nice observation on the Laporta algorithm: The integrals **A**, **C** without any numerator are of dimension $1/q^4$ (the fractional part is not important here) and thus power-counting finite, and they remain so with numerator $p_i \cdot q$. In our programmes we had not substituted explicit results for the master integrals till the very end, but rather kept the rational coefficient function that the Gaussian elimination runs up in the corresponding columns. Finiteness of **A**, **C** with any of the aforementioned numerators constrains the leading orders of the master integrals. The **B**, **D** graphs without a numerator are divergent due to their triangle subgraphs like e.g. the **BE** constellation at three loops. With numerator p_1^2 or p_2^2 **D** becomes finite, on **B** we may use any $p_i \cdot q$, $i \in \{1 \dots 4\}$.

The system of equations is strong enough to determine

$$\begin{aligned}
M_{35, 11111111} &= \frac{1}{(4\pi)^8} \left[\frac{\zeta(3)}{2\epsilon^2} + \left(\frac{\pi^4}{120} + \frac{11\zeta(3)}{2} \right) \frac{1}{\epsilon} + \left(-\frac{23\zeta(5)}{2} + \frac{11\pi^4}{120} + \frac{83\zeta(3)}{2} \right) \right. \\
&\quad \left. + \left(\frac{59\zeta(3)^2}{6} - \frac{2\pi^6}{63} - \frac{253\zeta(5)}{2} + \frac{83\pi^4}{120} + \frac{535\zeta(3)}{2} \right) \epsilon + \dots \right] \left(\frac{q^2}{\tilde{\mu}^2} \right)^{-4\epsilon}, \\
M_{36, 11111111} &= \frac{1}{(4\pi)^8} \left[\frac{5\zeta(5)}{\epsilon} + \left(-7\zeta(3)^2 + \frac{5\pi^6}{378} + 35\zeta(5) \right) + \dots \right] \left(\frac{q^2}{\tilde{\mu}^2} \right)^{-4\epsilon}, \\
NO_{\epsilon 11111111} &= \frac{1}{(4\pi)^6 q^2} \left[20\zeta(5) + \left(80\zeta(3)^2 + \frac{10\pi^6}{189} + 40\zeta(5) \right) \epsilon + \dots \right] \frac{(q^2)^{-4\epsilon}}{(\tilde{\mu}^2)^{-3\epsilon} (\mu^2)^{-\epsilon}}.
\end{aligned} \tag{89}$$

in exact agreement with the literature [16] upon conversion to the “G-scheme”. The given orders of the ϵ expansion of the masters are also all that our project required. The Laporta algorithm hence turned out to be self-sufficient.

Finiteness constraints on the level of the nine-propagator integral in the middle of Figure 12 determine one order less in M_{35} , M_{36} , and by finiteness of M_{35} , M_{36} constellations (we may introduce $p_i \cdot q / p_i^4$ on any line) we can only fix the first two terms in M_{35} while no condition on M_{36} ensues.

7 Conclusions

In this work we discussed the Konishi anomaly as an operator mixing problem in $\mathcal{N} = 4$ SYM with gauge group $SU(N)$. Up to normalisation the two operators that mix are the unintegrated chiral superpotential B and Yang-Mills action F of the $\mathcal{N} = 1$ superfield formulation of the theory. Both are not finite on their own, but the linear combination $O = F - 4gB$ is protected. The second mixture is fixed by the conformal properties of two-point functions, i.e. orthogonality to the protected operator and the form of its two-point function. We considered the first two non-trivial orders in perturbation theory using SSDR as a regulator and multiplicatively renormalised by Z-factors. In the result

$$K = Z_B B + \frac{gN}{32\pi^2} Z_F F \quad (90)$$

both renormalisation factors acquire singular higher loop corrections. Further, Z_F also receives non-vanishing finite corrections at higher loops. Our work fully confirms the general considerations about the singularities of the Z-factors put forward in [9] and extends the leading order perturbative analysis presented there. The descendant operator K has the anomalous dimension of the Konishi operator as required by supersymmetry.

In conclusion, in our framework the Konishi anomaly is not one-loop exact in contradiction to the comment after equation (2.110) in [8]. As an explanation we remark that higher loop mixing coefficients are usually scheme dependent. The approach quoted in [8] supposes a definite prescription for the renormalisation of the coupling constant appropriate to general $\mathcal{N} = 1$ theories, while our scheme is tailor-made for the conformal $\mathcal{N} = 4$ case. A general discussion in rigorous perturbation theory wide enough to reconcile the two contrary points of view is given in [17].

The technically hardest part of the project was to elaborate the four-loop two-point correlator $\langle \bar{B}F \rangle_{g^5}$, which we achieved by the Laporta algorithm, i.e. integration by parts paired with Gaussian elimination. We hope to separately publish the computer programmes developed to this end. Apart from the interest inherent to the renormalisation properties of the Konishi anomaly, the four-loop correlator is a vital piece of the calculation of the four-loop anomalous dimension of the operator K advocated in [18] along the lines of [14]; agreement with the existing result [19] would at the same time confirm the correctness of the method (thus the absence of a second “anomaly” in the supersymmetry variation of K) and further vindicate the aptitude of the thermodynamic Bethe ansatz [20] to describe “wrapping corrections” in the $\mathcal{N} = 4$ operator spectrum problem, i.e. the regime in which the dilatation generator in the sense of [21] becomes inapplicable because the loop order exceeds the spin chain length.

Acknowledgements

The author is deeply indebted to K. Chetyrkin for many discussions relating to the Laporta technique and for sharing results on the master integrals prior to publication. We are very grateful to A. Pak for an independent test of the numerator reduction by his own system. We thank S. Moch for stipulating our interest in the approach and K. Sibold for comments on the manuscript. Much of the computer work was done on a PC at the Spinoza Institute while the author was a member there. M. Segond pointed out *jaxodraw* as a convenient way of drawing Feynman graphs.

Appendix: Feynman rules and conventions

The $\mathcal{N} = 4$ SYM action formulated in terms of $\mathcal{N} = 1$ superfields has the form

$$\begin{aligned}
S_{\mathcal{N}=4} = & \int d^4x d^2\theta d^2\bar{\theta} \operatorname{Tr} (e^{gV} \bar{\Phi}_I e^{-gV} \Phi^I) \\
& + \left[\frac{g}{3!} \int d^4x_L d^2\theta \epsilon_{IJK} \operatorname{Tr} (\phi^I [\phi^J, \phi^K]) + c.c. \right] \\
& - \frac{1}{4g^2} \int d^4x_L d^2\theta \operatorname{Tr} (W^\alpha W_\alpha) \\
& - \frac{1}{\alpha} \int d^4x d^2\theta d^2\bar{\theta} \operatorname{Tr} \left(\left(-\frac{1}{4} D^2 V \right) \left(-\frac{1}{4} \bar{D}^2 V \right) \right) \\
& + \int d^4x d^2\theta d^2\bar{\theta} \operatorname{Tr} \left((b + \bar{b}) L_{\frac{g}{2} V} \left((c + \bar{c}) - \coth L_{\frac{g}{2} V} (c - \bar{c}) \right) \right).
\end{aligned} \tag{91}$$

The definition of the non-abelian field strength multiplet W_α is

$$W_\alpha = -\frac{i}{4} \bar{D}_{\dot{\alpha}} \bar{D}^{\dot{\alpha}} (e^{gV} D_\alpha e^{-gV}). \tag{92}$$

We choose Fermi-Feynman gauge by putting $\alpha = 1$: the quadratic part of the YM action becomes $+1/2 \int V \square V$. The action has the cubic and quartic YM self-interaction vertices

$$+ \frac{g}{4} \int d^4x d^2\theta d^2\bar{\theta} \operatorname{Tr} \left([V, (D^\alpha V)] \left(-\frac{1}{4} \bar{D}_{\dot{\alpha}} \bar{D}^{\dot{\alpha}} D_\alpha V \right) \right) \tag{93}$$

$$- \frac{g^2}{12} \int d^4x d^2\theta d^2\bar{\theta} \operatorname{Tr} \left([V, (D^\alpha V)] \left[V, \left(-\frac{1}{4} \bar{D}_{\dot{\alpha}} \bar{D}^{\dot{\alpha}} D_\alpha V \right) \right] \right) \tag{94}$$

$$+ \frac{g^2}{16} \int d^4x d^2\theta d^2\bar{\theta} \operatorname{Tr} \left([V, (D^\alpha V)] \left(-\frac{1}{4} \bar{D}_{\dot{\alpha}} \bar{D}^{\dot{\alpha}} [V, (D_\alpha V)] \right) \right). \tag{95}$$

For completeness, the first two ghost vertices are

$$+ \frac{g}{2} \int d^4x d^2\theta d^2\bar{\theta} \operatorname{Tr} \left((b + \bar{b}) [V, (c + \bar{c})] \right) \tag{96}$$

$$- \frac{g^2}{12} \int d^4x d^2\theta d^2\bar{\theta} \operatorname{Tr} \left([(b + \bar{b}), V] [V, (c - \bar{c})] \right) \tag{97}$$

W.r.t. the definitions in [3], the entire YM and ghost part of the action changes sign, which is related to the spinor convention in the integration measure. In addition, $V \leftrightarrow -V$. The ghost propagators are

$$\langle \bar{c}(2) b(1) \rangle = \langle \bar{b}(2) c(1) \rangle = \langle \bar{\phi}(2) \phi(1) \rangle. \tag{98}$$

In Minkowski space the action (91) appears in a weight factor e^{iS} under the path integral. Instead of putting the i onto each vertex we Wick rotate before determining the Feynman rules and proceed in positive Euclidean signature.

The superfields carry an adjoint representation of the gauge group $SU(N)$, and the (Hermitean) generators and the structure constants satisfy the relations

$$\operatorname{Tr}(T^a T^b) = \delta^{ab}, \quad f^{abc} f^{abd} = 2N \delta^{cd}. \tag{99}$$

The spinor convention is:

$$\psi^\alpha = \epsilon^{\alpha\beta} \psi_\beta, \quad \psi_\alpha = \epsilon_{\alpha\beta} \psi^\beta, \quad \epsilon_{12} = 1, \quad \epsilon_{\alpha\beta} \epsilon^{\beta\gamma} = \delta_\alpha^\gamma \tag{100}$$

and exactly the same with dotted indices. Complex conjugation replaces an undotted by a dotted index and vice versa; however, it does not exchange lower and upper position.

The 2×2 sigma matrices are Hermitian. The $\tilde{\sigma}$ matrix is obtained from $\sigma_{\alpha\dot{\alpha}}$ by raising of both indices as defined by the last equation. They satisfy the following relations:

$$\begin{aligned}
\sigma^\mu \tilde{\sigma}^\nu &= \eta^{\mu\nu} - i\sigma^{\mu\nu}, & \tilde{\sigma}^\mu \sigma^\nu &= \eta^{\mu\nu} - i\tilde{\sigma}^{\mu\nu}, \\
(\sigma^\mu)_{\alpha\dot{\alpha}} (\tilde{\sigma}_\mu)^{\dot{\beta}\beta} &= 2\delta_\alpha^\beta \delta_{\dot{\alpha}}^{\dot{\beta}}, & (\sigma^\mu)_{\alpha\dot{\alpha}} (\tilde{\sigma}_\mu)^{\dot{\alpha}\alpha} &= 2\delta_\nu^\mu.
\end{aligned} \tag{101}$$

The partial spinor derivative obeys

$$\partial_\alpha \theta^\beta = \delta_\alpha^\beta, \quad \bar{\partial}_{\dot{\alpha}} \bar{\theta}^{\dot{\beta}} = \delta_{\dot{\alpha}}^{\dot{\beta}}. \quad (102)$$

Derivatives with upper indices are defined by the raising rule given in the first equation in (100) and the same with dotted indices. (In case of doubt w.r.t. to signs the best strategy is always to put the indices into standard position: up on spinors and down on derivatives.) The superspace covariant derivatives are

$$D_\alpha = \partial_\alpha + i\bar{\theta}^{\dot{\alpha}}\partial_{\alpha\dot{\alpha}}, \quad \bar{D}_{\dot{\alpha}} = -\partial_{\dot{\alpha}} - i\theta^\alpha\partial_{\alpha\dot{\alpha}}, \quad \partial_{\alpha\dot{\alpha}} = \partial_\mu(\sigma^\mu)_{\alpha\dot{\alpha}}. \quad (103)$$

We define the square of a spinor without any weight factor:

$$\theta^2 = \theta^\alpha\theta_\alpha, \quad \bar{\theta}^2 = \bar{\theta}_{\dot{\alpha}}\bar{\theta}^{\dot{\alpha}}, \quad (104)$$

and similarly for the product of two different spinors. Consequently,

$$D.D\theta^2 \equiv -\frac{1}{4}D^2\theta^2 = 1, \quad \bar{D}.\bar{D}\bar{\theta}^2 \equiv -\frac{1}{4}\bar{D}^2\bar{\theta}^2 = 1. \quad (105)$$

Under the x -integral we may thus identify

$$d^2\theta = D.D, \quad d^2\bar{\theta} = \bar{D}.\bar{D}. \quad (106)$$

SSDR means to modify the dimension of space, but to leave the two-component spinor algebra untouched. In $D = 4 - 2\epsilon$ the basic propagator Π_{ij} (37) becomes

$$\Pi_{ij} = -\frac{\Gamma(1-\epsilon)\pi^\epsilon}{4\pi^2(x_{ij}^2)^{(1-\epsilon)}}\delta(\theta_{ij})\delta(\bar{\theta}_{ij}). \quad (107)$$

The propagators carry Kronecker delta symbols for colour and flavour indices, of course. We do not usually write these in order to unclutter the notation.

References

- [1] E. D'Hoker, D. Z. Freedman and W. Skiba, “Field theory tests for correlators in the AdS/CFT correspondence,” *Phys. Rev. D* **59** (1999) 045008 [arXiv:hep-th/9807098]; B. Eden, P. S. Howe and P. C. West, “Nilpotent invariants in $N = 4$ SYM,” *Phys. Lett. B* **463** (1999) 19 [arXiv:hep-th/9905085];
- [2] S. Penati, A. Santambrogio and D. Zanon, “Two-point functions of chiral operators in $N = 4$ SYM at order g^{**4} ,” *JHEP* **9912** (1999) 006 [arXiv:hep-th/9910197], “More on correlators and contact terms in $N = 4$ SYM at order g^{**4} ,” *Nucl. Phys. B* **593** (2001) 651 [arXiv:hep-th/0005223].
- [3] S. J. Gates, M. T. Grisaru, M. Rocek and W. Siegel, “Superspace, or one thousand and one lessons in supersymmetry,” *Front. Phys.* **58** (1983) 1 [arXiv:hep-th/0108200].
- [4] D. Anselmi, D. Z. Freedman, M. T. Grisaru and A. A. Johansen, “Universality of the operator product expansions of SCFT(4),” *Phys. Lett. B* **394** (1997) 329 [arXiv:hep-th/9608125]; B. Eden, C. Schubert and E. Sokatchev, “Three-loop four-point correlator in $N = 4$ SYM,” *Phys. Lett. B* **482** (2000) 309 [arXiv:hep-th/0003096]; M. Bianchi, S. Kovacs, G. Rossi and Y. S. Stanev, “Anomalous dimensions in $N = 4$ SYM theory at order g^{**4} ,” *Nucl. Phys. B* **584** (2000) 216 [arXiv:hep-th/0003203].
- [5] T. E. Clark, O. Piguet and K. Sibold, “Supercurrents, renormalization and anomalies,” *Nucl. Phys. B* **143** (1978) 445; “The absence of radiative corrections to the axial current anomaly in supersymmetric QED,” *Nucl. Phys. B* **159** (1979) 1.
- [6] K. Konishi, “Anomalous supersymmetry transformation of some composite operators in sqcd,” *Phys. Lett. B* **135** (1984) 439.

- [7] M. T. Grisaru, B. Milewski and D. Zanon, “The supercurrent and the Adler-Bardeen theorem,” Nucl. Phys. B **266** (1986) 589; M. A. Shifman and A. I. Vainshtein, “Solution of the anomaly puzzle in SUSY gauge theories and the Wilson operator expansion,” Nucl. Phys. B **277** (1986) 456 [Sov. Phys. JETP **64** (1986 ZETFA,91,723-744.1986) 428]; J. Erlich and D. Z. Freedman, “Conformal symmetry and the chiral anomaly,” Phys. Rev. D **55** (1997) 6522 [arXiv:hep-th/9611133]; D. Anselmi, M. T. Grisaru and A. Johansen, “A critical behaviour of anomalous currents, electromagnetic universality and CFT_4 ,” Nucl. Phys. B **491** (1997) 221 [arXiv:hep-th/9601023].
- [8] M. A. Shifman and A. I. Vainshtein, “Instantons versus supersymmetry: Fifteen years later,” arXiv:hep-th/9902018.
- [9] B. Eden, C. Jarczak, E. Sokatchev and Y. S. Stanev, “Operator mixing in $N = 4$ SYM: The Konishi anomaly revisited,” Nucl. Phys. B **722** (2005) 119 [arXiv:hep-th/0501077].
- [10] W. Siegel, “Supersymmetric dimensional regularization via dimensional reduction,” Phys. Lett. B **84** (1979) 193.
- [11] W. Siegel, “Inconsistency of supersymmetric dimensional regularization,” Phys. Lett. B **94** (1980) 37.
- [12] G. ’t Hooft and M. J. G. Veltman, “Regularization and renormalization of gauge fields,” Nucl. Phys. B **44** (1972) 189; K. G. Chetyrkin, A. L. Kataev and F. V. Tkachov, “New approach to evaluation of multiloop Feynman integrals: The Gegenbauer polynomial x space technique,” Nucl. Phys. B **174** (1980) 345, “Integration by parts: The algorithm to calculate beta functions in 4 loops,” Nucl. Phys. B **192** (1981) 159; F. V. Tkachov, “A theorem on analytical calculability of four loop renormalization group functions,” Phys. Lett. B **100** (1981) 65; D. I. Kazakov, “The method of uniqueness, a new powerful technique for multiloop calculations,” Phys. Lett. B **133** (1983) 406; S. A. Larin, F. V. Tkachov and J. A. M. Vermaseren, “The Form version of Mincer,” NIKHEF-H-91-18.
- [13] S. Laporta, “High-precision calculation of multi-loop Feynman integrals by difference equations,” Int. J. Mod. Phys. A **15** (2000) 5087 [arXiv:hep-ph/0102033].
- [14] B. Eden, C. Jarczak and E. Sokatchev, “A three-loop test of the dilatation operator in $N = 4$ SYM,” Nucl. Phys. B **712** (2005) 157 [arXiv:hep-th/0409009].
- [15] D. I. Kazakov, “Analytical methods for multiloop calculations: Two lectures on the method of uniqueness,” JINR-E2-84-410; D. J. Broadhurst, “Exploiting the 1.440 fold symmetry of the master two loop diagram,” Z. Phys. C **32** (1986) 249.
- [16] P. A. Baikov and K. G. Chetyrkin, in preparation.
- [17] E. Kraus, C. Rupp and K. Sibold, “Supersymmetric Yang-Mills theories with local coupling: The supersymmetric gauge,” Nucl. Phys. B **661** (2003) 83 [arXiv:hep-th/0212064]; “Conformal transformation properties of the supercurrent III: Nonabelian gauge theories with local coupling,” unpublished.
- [18] B. Eden, “Boxing with Konishi,” arXiv:0712.3513 [hep-th].
- [19] Z. Bajnok and R. A. Janik, “Four-loop perturbative Konishi from strings and finite size effects for multiparticle states,” Nucl. Phys. B **807** (2009) 625 [arXiv:0807.0399 [hep-th]]; F. Fiamberti, A. Santambrogio, C. Sieg and D. Zanon, “Anomalous dimension with wrapping at four loops in $N=4$ SYM,” Nucl. Phys. B **805** (2008) 231 [arXiv:0806.2095 [hep-th]]; V. N. Velizhanin, “The four-loop Konishi in $N=4$ SYM,” arXiv:0808.3832 [hep-th].
- [20] G. Arutyunov and S. Frolov, “On string S-matrix, bound states and TBA,” JHEP **0712** (2007) 024 [arXiv:0710.1568 [hep-th]]; N. Gromov, V. Kazakov and P. Vieira, “Exact Spectrum of Anomalous Dimensions of Planar $N=4$ Supersymmetric Yang-Mills Theory,” Phys. Rev. Lett. **103** (2009) 131601 [arXiv:0901.3753 [hep-th]]; D. Bombardelli, D. Fioravanti and R. Tateo, “Thermodynamic Bethe Ansatz for planar AdS/CFT: A Proposal”, J. Phys. A **42** (2009) 375401 [arXiv:0902.3930 [hep-th]]; N. Gromov, V. Kazakov, A. Kozak and P. Vieira, “Exact Spectrum of Anomalous Dimensions of Planar $N = 4$ Supersymmetric Yang-Mills Theory: TBA and excited states,” Lett. Math.

- Phys. **91** (2010) 265 [arXiv:0902.4458 [hep-th]]; G. Arutyunov and S. Frolov, “Thermodynamic Bethe ansatz for the $\text{AdS}_5 \times S^5$ mirror model,” JHEP **0905** (2009) 068 [arXiv:0903.0141 [hep-th]].
- [21] N. Beisert, C. Kristjansen and M. Staudacher, “The dilatation operator of $N = 4$ super Yang-Mills theory,” Nucl. Phys. B **664** (2003) 131 [arXiv:hep-th/0303060].